


A graphic of two stylized lungs in blue outline, positioned on either side of the title text.

# PULiMA

Series-MA

## BANGLADESH GUIDELINES FOR CHRONIC OBSTRUCTIVE PULMONARY DISEASE



**Mohammad Rashidul Hassan**





**BANGLADESH GUIDELINES FOR CHRONIC  
OBSTRUCTIVE PULMONARY DISEASE**

**MOHAMMAD RASHIDUL HASSAN**



**Author:**

Mohammad Rashidul Hassan

**Designed and Developed by:**

Ingenious Health Care Limited

41 Ring Road (2nd Floor), Mohammadpur, Dhaka-1207

Corresponding: E-mail: mrhassan07@gmail.com

Dhaka, Bangladesh

**Distributed by:**

- **Ingenious Pulmo-FIT**
- **Evidence Based Clinical Practice Society of Bangladesh (EBCPSB)**

[www.bdlungfoundation.org](http://www.bdlungfoundation.org)

Tel: +88 01911-792760

**First Edition:**

September, 2020

**Printing Courtesy:**

(As a service to respiratory medicine)

**Beximco Pharmaceuticals Ltd.**



**Price: Free**

*Contents of this book, whole or in part can be reproduced for research, academic or educational purpose. Acknowledgement to the author will be highly appreciated. No part of this book can be reproduced for commercial purpose as per copyright rules.*

# PREFACE

## PREFACE

By the grace of Almighty Allah, we are able to publish our own Bangladeshi COPD guidelines, which is evidence based and not a true copy of any other guidelines. We tried to focus that pin point diagnosis is important for a COPD patient as COPD is not a single disease, rather it's a cluster of diseases.

Spirometry assessment is important for diagnosis of severity and staging on the basis of  $FEV_1$ . But post-bronchodilator  $FEV_1/FVC$  ratio  $<0.7$  is not a reliable tool for diagnosis of COPD as a number of patient including under 50 years with COPD and some may have associated mixed Lung parenchymal fibrosis and/or multiple Bullae which may cause post-bronchodilator  $FEV_1/FVC$  ratio  $>0.7$ .

We intentionally avoided “ABCD” assessment tool of GOLD guidelines as many co-existed pulmonary diseases may have same symptoms and affects CAT Score and/or breathlessness of COPD by the Modified British Medical Research Council (mMRC) Questionnaire. We tried to explain that simple COPD diagnosis and spirometry assessment is not sufficient for a management plan of COPD. The combination of airflow limitation along with clinical parameters and investigation tools makes it clear to find out clusters of diseases present in an individual with COPD. All diagnostic point should be evaluated for diagnosis, staging and classification of disease. Our assessment tool acknowledges Disease diagnosis, rate of exacerbations per year and  $FEV_1$  which will determine treatment plan for an individualized COPD patient. It will highlight the importance of patients' cluster of disease and exacerbation risks in patient with COPD. Spirometry remains an important key in the diagnosis of  $FEV_1$  staging, prognostication and treatment with non-pharmacological therapies.

GOLD 2020 revised guidelines stated that the blood eosinophil count as a biomarker and as a guide to prescribe and use inhaled corticosteroids for the prevention of exacerbations. But it is only one parameter for diagnosis of asthma in COPD patient and they used in group D patient only.

We emphasize that asthma and COPD are different disorders although they share some traits and features (e.g., eosinophilia, some degree of reversibility). It is now clear that like asthma in a COPD patient, a physician should try to find out other diseases like bronchiectasis, GERD and OSA. They are also not less common than asthma and they are developing during the progress of COPD.

Meticulous observation and follow up is required to establish this concept of COPD management and patient will get more dependable management from care giver throughout the world.

We are grateful to our committee for their support to make this effort successful.

### **Mohammad Rashidul Hassan**

Chairman, Ingenious Health Care Limited  
Professor of Respiratory Medicine  
Anwer Khan Modern Medical College, Dhaka  
Bangladesh

# Bangladesh COPD Guidelines Committee

- Editor-in-Chief** : **Mohammad Rashidul Hassan**  
Ingenious Health Care Limited, Dhaka, Bangladesh
- Editorial Board** : **Abu NGA Khan**  
Nyp Morgan Stanley Childrens Hospital, New York, USA
- Chief Reviewer** : **GM Monsur Habib**  
The University of Edinburgh (UK)
- Rawshan Aara Khanam**  
United Hospital Dhaka, Bangladesh
- Arman Faisal**  
Ingenious Health Care Limited, Dhaka, Bangladesh
- Pulak Kumar Dey**  
NIDCH, Dhaka, Bangladesh
- Shamima Akter**  
NIDCH, Dhaka, Bangladesh
- Jalal Mohsin Uddin**  
NIDCH, Dhaka
- Raj Kumar**  
Vallabhbhai Patel Chest Institute, New Delhi, India
- SM Rowshan Alam**  
RMCH, Rangpur, Bangladesh
- S.M Sanayet Bin Mustafiz**  
United Hospital, Dhaka, Bangladesh
- Mohammad Towfique Hasan**  
Sheikh Hasina Medical College, Tangail, Bangladesh
- Taibee Jhumara**  
Dh. Rinbudhoo Health Center, Maldiv
- Md. Abdulla-hel Kafee**  
Kurmitola General Hospital, Dhaka, Bangladesh
- Muhammad Murtaza Khair**  
NIDCH, Dhaka, Bangladesh
- AKM Akramul Haque**  
NIDCH, Dhaka
- Md. Zakir Hossain Sarker**  
NIDCH, Dhaka, Bangladesh
- Abdullah-Hel-Kafi Khan**  
Anwer Khan Modern Medical College, Dhaka, Bangladesh

## Past and Present Contributing Panel and Advisory Board :

|   |  |
|---|--|
| <p><b>Md Ali Hossain</b><br/>NIDCH, Dhaka, Bangladesh</p> <p><b>Asif Mujtaba Mahmud</b><br/>Asgar Ali Hospital, Dhaka, Bangladesh</p> <p><b>Md. Mostafizur Rahman</b><br/>Modern Medical College, Dhaka, Bangladesh</p> <p><b>Mohammad Mohiuddin Ahmad</b><br/>DMC, Dhaka, Bangladesh</p> <p><b>Hena Khatun</b><br/>DMCH, Dhaka, Bangladesh</p> <p><b>Mohammad Abdus Shakur Khan</b><br/>NIDCH, Dhaka, Bangladesh</p> <p><b>Samir C Majumder</b><br/>RMC, Rajshahi, Bangladesh</p> <p><b>Rahmatul Bari</b><br/>Dhaka, Bangladesh</p> <p><b>B.K Bose</b><br/>Dinajpur, Bangladesh</p> <p><b>Md. Delwar Hossain</b><br/>BIRDEM, Dhaka, Bangladesh</p> | <p><b>Sk Shahinur Hossain</b><br/>Dhaka, Bangladesh</p> <p><b>Dewan Azmal Hussain</b><br/>NIDCH, Dhaka, Bangladesh</p> <p><b>Md. Ferdous Wahid</b><br/>NIDCH, Dhaka, Bangladesh</p> <p><b>Ashrafal Alam Khan, NIDCH,</b><br/>Dhaka, Bangladesh</p> <p><b>Kazi Fawzia Afreeen</b><br/>Dhaka, Bangladesh</p> <p><b>AKM Shamsul Huq</b><br/>Dhaka, Bangladesh</p> <p><b>Md. Ruhul Amin</b><br/>Dhaka, Bangladesh</p> <p><b>ARM Luthful Kabir</b><br/>Dhaka, Bangladesh</p> <p><b>Muhammad Rafiqul Alam</b><br/>BSMMU, Dhaka, Bangladesh</p> |
|---|--|

## Past and Present Facilitators :

|   |   |
|---|---|
| <p><b>Md. Abdul Qayyum</b><br/>Dhaka, Bangladesh</p> <p><b>Mohammad Delwar Hossain</b><br/>Sylhet, Bangladesh</p> <p><b>ASM Areef Ahsan</b><br/>BIRDEM, Dhaka, Bangladesh</p> <p><b>Sheikh AHM Mesbahul Islam</b><br/>Dhaka, Bangladesh</p> <p><b>Adnan Yusuf Choudhury</b><br/>DMC, Dhaka, Bangladesh</p> <p><b>S M Zakir Hossain</b><br/>Dhaka, Bangladesh</p> <p><b>Md. Shahan</b><br/>Dhaka, Bangladesh</p> | <p><b>Mustafijur Rahman</b><br/>Dhaka, Bangladesh</p> <p><b>Md. Safiul Islam</b><br/>Dhaka, Bangladesh</p> <p><b>Md. Masudur Rahman</b><br/>Dhaka, Bangladesh</p> <p><b>Masum Ahmed</b><br/>Barisal, Bangladesh</p> <p><b>Md. Mamunur Rashid</b><br/>Dhaka, Bangladesh</p> <p><b>Miraz Mahmud</b><br/>Dhaka, Bangladesh</p> <p><b>Md. Shahjada Tabrez</b><br/>Dhaka, Bangladesh</p> |
|---|---|

## Acknowledgement :

|  |   |
|--|---|
| <p><b>Mohammad Jubayer Hassan Tamim</b><br/><b>Foysal Ahmed</b><br/><b>Refat Mahmood</b></p> | <p>Beximco Pharmaceuticals Ltd<br/>Incepta Pharmaceuticals Ltd<br/>Eskayef Pharmaceuticals Ltd<br/>UniMed &amp; UniHealth Mfg. Ltd<br/>Healthcare Pharmaceuticals<br/>ACME Laboratories Ltd</p> |
|--|---|

# INDEX

## LIST OF TABLE:

|   |    |
|---|----|
| Table 1: Mnemonic meaning of 'ABG' & 'PTO'                            | 30 |
| Table 2: Severity of COPD on the basis of FEV <sub>1</sub> GOLD 2020. | 35 |
| Table 3: Modified Medical Research Council (mMRC) Scale               | 40 |
| Table 4: Modified Medical Research council scale in Bangla            | 40 |
| Table 5: Calculation for PaO <sub>2</sub> assessment                  | 45 |
| Table 6: Diagnosis of COPD-A:   | 49 |
| Table 7: Diagnosis of COPD-B  | 51 |
| Table 8: Diagnosis of COPD-G  | 54 |
| Table 9: HRD score Uploaded by Rachel Mary Thomson                    | 55 |
| Table 10: Diagnosis of COPD-O   | 58 |
| Table 11: Epworth Sleepiness Scale                                    | 59 |
| Table 12: STOP-BANG Questionnaire                                     | 60 |
| Table 13: Cut-off values of BNP and NTproBN                           | 64 |
| Table 14: Diagnosis of COPD and Heart Failure                         | 65 |
| Table 15: Criteria for Diagnosis of the Frailty Syndrome              | 67 |
| Table 16: FRAIL SYNDROME: QUESTIONNAIRE BASED DIAGNOSIS               | 67 |
| Table 17: Treatments of Frailty Syndrome                              | 68 |
| Table 18: Non-pharmacological therapy for all COPD                    | 74 |
| Table 19: Non-pharmacological therapy for specific COPD               | 74 |
| Table 20: Degree of Physical Activity                                 | 78 |
| Table 21: Components of Physical Activity                             | 80 |
| Table 22: Pharmacotherapy in COPD                                     | 86 |
| Table 23: Pharmacotherapy for COPD-B                                  | 87 |
| Table 24: Pharmacotherapy of COPD-G                                   | 87 |
| Table 25: Pharmacotherapy of COPD - O & S                             | 88 |
| Table 26: Pharmacotherapy of COPD - U                                 | 89 |
| Table 27: Classification of benzodiazepine                            | 90 |
| Table 28: Diagnostic features of Pneumonia                            | 92 |
| Table 29: Diagnostic features of Pneumothorax                         | 92 |
| Table 30: Diagnostic features of Pleural effusion                     | 93 |
| Table 31: Features of Pulmonary embolism                              | 93 |
| Table 32: Features of Pulmonary edema                                 | 93 |
| Table 33: Identifying Cardiac arrhythmias                             | 94 |
| Table 34: Cut off Values of CRP for COPD needs antibiotics            | 94 |



# INDEX

## LIST OF FIGURE:

|   |    |
|---|----|
| Figure 1: Different phenotypes of COPD                              | 19 |
| Figure 2: Clusters of COPD phenotype                                | 22 |
| Figure 3: Non-proportional Phenotypes of COPD                       | 23 |
| Figure 4: Proportional Venn Diagram of COPD                         | 24 |
| Figure 5: Disease trajectory of COPD                                | 25 |
| Figure 6: Risk of occupational exposure for COPD                    | 27 |
| Figure 7: Two distinct infection cycles in COPD                     | 29 |
| Figure 8: A Journey of Stable to Unstable COPD                      | 33 |
| Figure 9: Coexisting diseases and comorbidities in COPD.            | 34 |
| Figure 10: Algorithm of diagnosis of COPD and Coexisting disease    | 37 |
| Figure 11: Algorithm for Management of COPD                         | 38 |
| Figure 12: CAT SCORE  | 39 |
| Figure 13: CAT SCORE (Bangla Version)                               | 39 |
| Figure 14: Comparison of different flow-volume curves               | 42 |
| Figure 15: Causes of breathlessness                                 | 42 |
| Figure 16: Oxyhemoglobin dissociation curve                         | 45 |
| Figure 17: Calculation of absolute ratio of improvement             | 50 |
| Figure 18: CT Scan at upper and lower level of chest                | 56 |
| Figure 19: STOP-Bang questionnaire                                  | 60 |
| Figure 20: Self-reported questionnaire tool for frailty             | 67 |
| Figure 21: Images of combined pulmonary fibrosis and emphysema      | 71 |
| Figure 22: Algorithm of COPD MANAGEMENT                             | 73 |
| Figure 23: Patho-physiology of exercise limitation                  | 76 |
| Figure 24: COPD's vicious circle (reproduced from).                 | 77 |
| Figure 25: physical activity in urban and rural of Bangladesh.      | 79 |
| Figure 26: Assessment of Physical Activity in Pulmo-Fit             | 81 |
| Figure 27: Mechanism of ECMO  | 83 |
| Figure 28: PCT in different infections                              | 95 |
| Figure 29: PCT algorithm with respiratory tract infections in A & E | 96 |
| Figure 30: PCT algorithm with sepsis in the ICU                     | 97 |

## CONTENTS

|   |    |
|---|----|
| BANGLADESH GUIDELINES FOR COPD                          | 1  |
| PREFACE   | 3  |
| Bangladesh COPD Guidelines Committee                    | 4  |
| Past and Present Contributing Panel and Advisory Board: | 5  |
| Acknowledgement :                                       | 5  |
| List of Table   | 6  |
| List of Figure:   | 7  |
| <b>CONTENTS</b>   | 8  |
| List of Abbreviations:                                  | 12 |
| <b>A BACKGROUND</b>                                     | 14 |
| <b>B Executive Summary:</b>                             | 15 |
| <b>1 CHAPTER 1: INTRODUCTION</b>                        | 18 |
| 1.1 Bangladesh perspective:                             | 18 |
| 1.2 Management strategy of COPD:                        | 20 |
| <b>2 CHAPTER 2: METHODOLOGY</b>                         | 21 |
| 2.1 Guideline development:                              | 21 |
| <b>3 CHAPTER 3: DEFINITION</b>                          | 22 |
| 3.1 Definitions in short                                | 22 |
| 3.2 Clinical definition of COPD:                        | 23 |
| 3.2.1 Pre-COPD or Early COPD:                           | 26 |
| <b>4 CHAPTER 4: DEVELOPMENT AND PROGRESSION of COPD</b> | 27 |
| <b>5 CHAPTER 5: OVERVIEW</b>                            | 29 |
| <b>6 CHAPTER 6: DEF CLASSIFICATION AND STAGING</b>      | 32 |
| 6.1 Distinct pathophysiological features:               | 33 |
| 6.2 Cardio-Pulmonary Exercise Test (CPET)               | 34 |
| 6.3 SEVERITY OF COPD:                                   | 35 |
| <b>7 CHAPTER 7: DIAGNOSIS</b>                           | 36 |
| 7.1 Summary notes:                                      | 36 |

|           |  |           |
|-----------|--|-----------|
| <b>8</b>  | <b>CHAPTER 8: INVESTIGATIONS</b>                               | <b>41</b> |
|           | 8.1 CBC:   | 41        |
|           | 8.2 Chest X-Ray  | 41        |
|           | 8.3 Sputum examination   | 41        |
|           | 8.4 Spirometry   | 41        |
|           | 8.5 Exercise testing   | 42        |
|           | 8.5.1 Components of Modified CPET:                             | 43        |
|           | 8.6 Fractional exhaled Nitric Oxide (FeNO)                     | 44        |
|           | What is FeNO breathing test?                                   | 44        |
|           | Advantages of FeNO:  | 44        |
|           | Japanese Recommendation:                                       | 44        |
|           | 8.7 Saturation of pulsatile O <sub>2</sub> (SpO <sub>2</sub> ) | 44        |
|           | 8.8 Arterial blood gas measurement                             | 45        |
|           | 8.9 Biochemistry   | 46        |
|           | 8.10 E.C.G and Echo ?  | 46        |
|           | Usefulness of Echocardiography in COPD:                        | 47        |
|           | 8.11 High resolution computed tomography                       | 47        |
|           | 8.12 Sleep studies   | 47        |
| <b>9</b>  | <b>CHAPTER 9: SUB-CLASSIFICATION OF COPD</b>                   | <b>48</b> |
|           | 9.1 COPD with Asthma: COPD-A                                   | 48        |
|           | 9.2 COPD and Bronchiectasis: COPD-B                            | 50        |
|           | 9.3 COPD and GERD: COPD-G                                      | 53        |
|           | 9.4 COPD and Obstructive Sleep Apnea (OSA):                    |           |
|           | COPD-O   | 56        |
|           | 9.4.1 Sleep and Breathing                                      | 57        |
|           | 9.4.2 Nocturnal Oxygen Desaturation                            | 57        |
|           | 9.4.3 Screen for Obstructive Sleep Apnea (OSA):                | 59        |
|           | 9.5 Mixed Disorder:  | 60        |
| <b>10</b> | <b>CHAPTER 10: COMMON PROBLEM FOR COPD</b>                     | <b>61</b> |
|           | 10.1 Major triggers, common to all                             |           |
|           | COPD patients:   | 61        |
|           | 10.1.1 Air Pollution:  | 61        |
|           | 10.1.2 Temperature   | 61        |
| <b>11</b> | <b>CHAPTER 11: COMORBIDITIES IN COPD</b>                       | <b>62</b> |
|           | 11.1 Comorbidities:  | 62        |
|           | 11.2 Cardiac Disease   | 62        |
|           | 11.3 Heart Failure   | 63        |
|           | 11.3.1 Role of BNP and NT-proBNP in                            |           |
|           | clinical practice  | 63        |

# INDEX

|        |  |    |
|--------|--|----|
| 11.3.2 | Use of Cardio-selective Beta Blocker in COPD:          | 66 |
| 11.4   | Osteoporosis   | 66 |
| 11.5   | Frailty and COPD                                       | 66 |
| 11.5.1 | Management of Frailty:                                 | 68 |
| 11.6   | Falls in COPD  | 69 |
| 11.7   | Aspiration and COPD                                    | 69 |
| 11.7.1 | Eating strategies which may prevent dyspnea            | 70 |
| 11.8   | Lung cancer and chronic obstructive pulmonary disease: | 71 |
| 11.9   | Combined Pulmonary Fibrosis and Emphysema              | 71 |

|           |   |           |
|-----------|---|-----------|
| <b>12</b> | <b>CHAPTER 12: NON-PHARMACOLOGICAL MANAGEMENT</b>   | <b>74</b> |
| 12.1      | For All Patients of COPD:                           | 74        |
| 12.2      | Pulmonary Rehabilitation (PR):                      | 75        |
| 12.2.1    | Patho-physiology of exercise limitation:            | 75        |
| 12.2.2    | Integrative Care in COPD Pulmonary Rehabilitation   | 76        |
| 12.2.3    | Physical Activity                                   | 77        |
| 12.2.4    | Education and self-management                       | 81        |
| 12.3      | Non-Invasive Ventilation (NIV):                     | 81        |
| 12.3.1    | a. Acute NIV:                                       | 81        |
| 12.3.2    | b. Chronic NIV:                                     | 81        |
| 12.4      | Long Term Oxygen Therapy (LTOT):                    | 82        |
| 12.5      | Ambulatory Oxygen Therapy                           | 82        |
| 12.6      | Short-Burst Oxygen Therapy                          | 82        |
| 12.7      | Extracorporeal Membrane Oxygenation (ECMO):         | 83        |
| 12.8      | Surgery   | 84        |
| 12.8.1    | Bullectomy  | 84        |
| 12.8.2    | Lung Volume Reduction and Endobronchial Procedures: | 84        |
| 12.8.3    | Lung Transplantation:                               | 85        |
| 12.9      | End of Life (EOL)Care:                              | 85        |

# INDEX

|           |   |            |
|-----------|---|------------|
| <b>13</b> | <b>CHAPTER 13: PHARMACOLOGICAL TREATMENT OF COPD</b>                | <b>86</b>  |
| 13.1      | Pharmacotherapy in COPD   | 86         |
| 13.2      | Treatment of anxiety and depression in COPD                         | 89         |
| <b>14</b> | <b>CHAPTER 14: MANAGEMENT OF EXACERBATIONS OF COPD (Plan ABCO):</b> | <b>91</b>  |
| 14.1      | Exacerbations of COPD   | 91         |
| 14.2      | COPD Exacerbations PLAN (ABCO):                                     | 94         |
| 14.3      | A. Antibiotics for treatment of Exacerbations:                      | 94         |
| 14.4      | systemic corticosteroids + antibiotics for exacerbation             | 98         |
| 14.5      | B. Inhaled bronchodilators in exacerbations                         | 98         |
| 14.5.1    | Effectiveness of Inhaled BD COPD exacerbations                      | 98         |
| 14.5.2    | C. Short course Systemic corticosteroids                            | 98         |
| 14.6      | O. Oxygen during exacerbations:                                     | 99         |
| 14.6.1    | Controlled oxygen therapy:  | 99         |
| 14.7      | Non-Invasive Ventilatory assistance:                                | 99         |
| <b>15</b> | <b>CHAPTER 15: COPD AND ASSESSMENT FOR SURGERY</b>                  | <b>100</b> |
| 15.1      | Pre-operative Assessment for COPD patient:                          | 100        |
| 15.1.1    | Lung function assessment for lung surgery                           | 100        |
| <b>16</b> | <b>CHAPTER 16. REGENERATIVE THERAPY</b>                             | <b>101</b> |
| 16.1      | A. Stem cell therapy for (COPD)                                     | 101        |
| 16.2      | B. Exosome in COPD :  | 101        |
| <b>17</b> | <b>CHAPTER 17: References:</b>                                      | <b>102</b> |

## LIST OF ABBREVIATIONS:

|        |   |
|--------|---|
| ABG    | Arterial Blood Gas                        |
| AF     | Atrial Fibrillation                       |
| BNP    | B-type natriuretic peptide                |
| CAT    | COPD Assessment Test                      |
| CBC    | Complete Blood Count                      |
| CHD    | Coronary Heart Disease                    |
| CHF    | Congestive Heart Failure                  |
| COPD   | Chronic Obstructive Pulmonary Disease.    |
| CPFE   | Combined Pulmonary Fibrosis and Emphysema |
| CRF    | Chronic Renal Failure;                    |
| CRD    | Chronic Respiratory Disease               |
| DEF    | Disease Based Classification              |
| ECMO   | Extra-Corporeal Membrane Oxygenation      |
| FeNO   | Fractional Exhaled Nitric Oxide           |
| GERD   | Gastro-Esophageal Reflux Disease          |
| GPAQ   | Global Physical Activity Questionnaire    |
| HIC    | High Income Countries                     |
| HR     | Heart Rate                                |
| HRCT   | High resolution computed tomography       |
| HRD    | Heart burn, Regurgitation, Dysphagia      |
| HTN    | Hypertension                              |
| ICS    | Inhaled Corticosteroid                    |
| LABA   | Long Acting Beta-2 Agonist                |
| LAMA   | Long-Acting Muscarinic Antagonists        |
| LLN    | Lower Limit of Normal                     |
| LMIC   | Low- and Middle-Income Countries          |
| LTOT   | Long Term Oxygen Therapy                  |
| LVRS   | Lung Volume Reduction Surgery             |
| M-CPET | Modified Cardio-Pulmonary Exercise Test   |
| MET    | Metabolic Equivalent Task                 |
| mMRC   | Modified Medical Research Council         |

# INDEX

|       |  |
|-------|--|
| MS    | Metabolic Syndrome                             |
| NCD   | Non-Communicable Diseases                      |
| NIV   | Non-Invasive Ventilation                       |
| OSA   | Obstructive Sleep Apnea                        |
| OSAS  | Obstructive Sleep Apnea Syndrome               |
| PA    | Physical Activity                              |
| PiA   | Physical inactivity                            |
| PHRHF | Pulmonary Hypertension and Right Heart Failure |
| PR    | Pulmonary Rehabilitation                       |
| PTO   | Patient may Turn Out                           |
| REM   | Rapid-Eye-Movement                             |
| VTE   | Venous Thrombo-Embolism                        |

## Disclaimer:

Recommendation of this guideline might not be successful in every situation. Clinical judgement, patient's preference, and overall practice protocol should be considered in every point of implementation of the guideline. The guideline echoes the best evidence-based data, recommendation, and guidance for the best care of COPD patients. Furthermore, updated information should be added once available because of rapidly budding evidence in medical science. As a part of continuously upgrading medical care, this guideline will be re-published once major alterations are available to imitate new data.

### A. BACKGROUND

The aim of this guideline is to provide an evidence-based deliverable, user-friendly guideline for Bangladeshi and physicians who are working in low- and middle-income countries (LMICS), as well as those who are interested to give more precise support to COPD patients throughout the world. We want to develop and spread a new strategy of COPD management.

Aim of this guideline firstly is to establish proper diagnosis of COPD, secondly, to diagnose coexisting diseases and all possible comorbidities. Thirdly, is to classify COPD severity on the basis of spirometry, i.e. Lung Function. This is based on  $FEV_1$  and is called COPD staging to understand patient's disease progression, by which severity of airflow limitation can be tracked. Fourthly, is to establish diagnosis whether patient has COPD Stable state, COPD with/ or without coexisting illnesses and with/ or without unstable COPD on the basis of frequency of exacerbations or hospitalization per year<sup>1</sup>. Much is now known about the characteristics of patients in the different COPD stages. Risk of exacerbations, hospitalization, and death of COPD patients depends more on coexisting illnesses and comorbidities than COPD staging<sup>2</sup>. For this reason, at an individual patient level,  $FEV_1$  may be an unreliable marker of the severity of breathlessness, exercise limitation, and health status impairment. Level of management depends on 3 basic assessments: Diagnosis of Diseases, Frequency of Exacerbations and Staging of COPD (DEF i.e. Disease Exacerbation and  $FEV_1$ ).



## B. Executive Summary:

- COPD is one of the progressively deteriorating burden in the health care system of Bangladesh.
- Evidence-based, appropriate, deliverable management guideline is the greatest standing in this field.
- Protocol of COPD management in High-Income Countries (HICs) might not be deliverable and applicable in Bangladesh as well as in LMICS.
- Aim of this guideline:
  - Provide fundamentals of accurate diagnosis of COPD
  - Identifying the coexisting diseases and co-morbidities
  - Spirometric classification of the severity disease (on FEV<sub>1</sub>)
  - Disease based classification (DEF) for ensuring optimal therapy.
  - Identifying COPD as:
    - Stable COPD
    - ± coexisting disease
    - ± unstable COPD (based on exacerbation or hospitalization rate)
- Risk of exacerbations, hospitalization, and death of COPD patients depends more on coexisting illnesses and comorbidities than conventional COPD staging.
- Despite sputum production being a common symptom, usually it is under-reported by most of the patients<sup>3</sup>.
- The main risk factor for COPD is tobacco smoking but other environmental exposures such as biomass fuel exposure and air pollution may contribute.
- Biological dust exposures include substances of microbial, plant or animal origin such as bacteria, fungi, allergens, endotoxins, peptidoglycans, glucans, pollens, and plant fibers.
- Lung function usually deteriorate after exacerbations and usually never return to previous state after each exacerbation in COPD,
- How fast the lung function will deteriorate, depends on the commencing risk factors of the disease.
- Coexisting and comorbid conditions of COPD upsurges the morbidity and mortality.
- Compared to smokers with COPD, nonsmokers with chronic airflow limitation have fewer symptoms, milder disease and lower burden of systemic inflammation, however, non-smokers have higher risk of pneumonia.
- The development of COPD is most likely a series of complex interactions between noxious airborne agents together with a genetic predisposition.
- FEV<sub>1</sub> may be an unreliable marker of the severity and symptom score marker

- The goals of effective COPD management are to:
  - Prevent disease progression by controlling Risk factors
  - Relieve symptoms by modern Pharmacological medicine
  - Improve exercise tolerance by Pulmonary Rehabilitation Program
  - Improve health status by controlling coexisting diseases and comorbidities
  - Prevent and treat complications by precise diagnosis of all components of COPD
  - Prevent and treat exacerbations by vaccination and avoiding air pollution, temperature changes and early detection and treating infection
  - Reduce mortality by reducing exacerbations, hospitalizations frail syndrome and by controlling coexisting diseases and comorbidities.
- Management plan depends on 3 basic parameters of COPD:
  - Diagnosis of Diseases, - (D)
  - Frequency of Exacerbations and (E)
  - Severity of COPD (FEV<sub>1</sub>).(F)
- Theme sentence of this guideline is: *"If Arterial Blood Gas (ABG) deteriorates, stable COPD Patient may Turn Out (PTO) to unstable COPD"*.
- Here actually 'A' means Asthma, 'B' means Bronchiectasis, G means Gastro-Esophageal Reflux Disease (GERD), 'P' means air Pollution, 'T' means Temperature changes and 'O' means Obstructive Sleep Apnea (OSA).
- Stable COPD: (COPD- S) i.e. Indolent COPD, having no coexisting lung disease with having one or less exacerbation per year and virtually no hospitalization for COPD
  - COPD with coexisting diseases: COPD-C
  - COPD with Asthma: COPD-A
  - COPD and Bronchiectasis: COPD-B
  - COPD and GERD: COPD-G
  - COPD and Obstructive Sleep Apnea (OSA): COPD-O
  - Unstable COPD: (COPD – U); COPD U means COPD patient having more than two exacerbations per year without hospitalization or one exacerbation with hospitalization per year.
- Any one of 15 subtypes of COPD C group may turn to COPD U if they develop frequent exacerbations or hospitalization in a year
- For precise diagnosis and sub-categorization (single or combinations) of COPD C, we need some investigations i.e. a good quality Chest X-ray, HRCT scan of chest (if possible), CBC, and total circulating eosinophil count, sputum cytology with differential count, Blood total IgE level and FeNO.

- Diagnosis part has Three Components:
  - Diagnosis of COPD
  - Diagnosis of Coexisting Diseases
  - Diagnosis of Comorbidities
- For the diagnosis of COPD, comorbidity and coexisting diseases certain investigations have paramount importance e.g. CBC, radiological examination of chest, sputum examination, exercise testing (modified CPET), FeNO, ABG, biochemistry values, ECG, and Sleep studies.
- Common comorbid conditions of COPD are, heart failure, osteoporosis, frailty syndrome, falls in COPD, aspiration, lung cancer, and combined fibrosis with emphysema.
- Management of COPD: for the proper management of COPD precise diagnosis and classification is mandatory. This guideline suggests DEF classification for specifying the management option. Thereafter, Non-pharmacological and pharmacological therapy should be determined (Table16).
- Non-pharmacological therapy includes, reducing exposure to risk factors including quitting smoking, avoidance of temperature fluctuation, Pulmonary Rehabilitation, increasing physical activities, developing skill for self-care, chronic NIV, long-term, ambulatory, and short burst oxygen therapy, ECMO, and surgical intervention.
- Management of comorbidity deserves paramount importance in the care of COPD, as such proper identification and appropriate intervention is statutory.
- Anxiety and depression are common in COPD and that should be addressed sensibly.
- COPD is punctuated by exacerbation, that's why an 'exacerbation management plan' should be pre-defined. In this guideline we used the acronym the plan as 'ABCO' (A-Antibiotic, B-Bronchodilator, C-Corticosteroid, and O-Oxygen)-Chapter
- Before the surgical intervention a precise assessment of lung function is mandatory which is described in chapter 15
- New epoch of the management of COPD is stem cell therapy which is discussed briefly in chapter 16.

# 1 CHAPTER 1: INTRODUCTION

According to WHO estimates, 65 million people have moderate to severe chronic obstructive pulmonary disease (COPD)<sup>4</sup>. COPD is currently the fourth leading cause of death in the world but is projected to be the 3rd leading cause of death by 2030.

More than 3.2 million people died of COPD in 2015<sup>5</sup>, which corresponds to 5% of all deaths globally. It is known that almost 90% of COPD deaths occur in LMICs<sup>6</sup>. At one time, COPD was more common in men, but because of increased tobacco use among women in HICs and the higher risk of exposure to indoor air pollution (such as biomass fuel used for cooking and heating) in LMICs, the disease now affects men and women almost equally<sup>7</sup>.

## 1.1 Bangladesh perspective:

According to Burden of Obstructive Lung Diseases Bangladesh (BOLD BD study 2007), in Bangladesh prevalence of COPD in total population of Bangladesh was 4.24% and estimated 6.0 million people were suffering from COPD<sup>8</sup>. The Prevalence in population of 40 years and above was found to be 21.58% with a total of 60,42,400 people. Majority (45.13%) of the COPD patients were suffering from GOLD spirometry classification of moderate stage of COPD. In general, males suffer more than females (62.47% vs. 37.26%). Prevalence in urban areas (61.52%) is higher than that of rural areas (38.48%). Interestingly, among the female COPD patients, these figures are almost equal (urban ~ 50.21% vs. rural 49.79%), indicating higher prevalence of the disease in village women<sup>9</sup>.

From these findings, it is obvious that COPD is a disease for the poorer people (~80% of total population) of Bangladesh. People of the productive age group (40-50 years) was the main victim (42%). The prevalence decreases with the increase of age (lowest in more than 80 years group – 2.5%). About 50% patients with COPD were illiterate.

Majority (63%) COPD patients were current smokers. Smoker group was extremely high in male COPD patients – about 88% were smokers, whereas only 20% of female COPD patients were smokers. We found a positive relation between gradual raise in COPD prevalence with increased pack-years of cigarette or equivalent smoking tobacco consumption. It showed that majority of smokers (80%) need only around 10 pack-years of smoking to develop the disease. Exposure to biomass gas (open stove/wood burn) appears to be a significant risk factor in COPD. More than 42% COPD patients are exposed to it. The figure was even higher among female COPD patients (more than 86%).




Figure 1: Different phenotypes of COPD

1. COPD
2. COPD with coexisting diseases
3. COPD with comorbidities
4. COPD with coexisting diseases and comorbidities
5. 6. 7. Some diseases without COPD but having other obstructive lung diseases like asthma
8. 9. Some diseases without obstructive lung disease

- The large Triangle represents total diagnosed and undiagnosed Obstructive Pulmonary Diseases.
- The circles represent the Nonproportional overlapping of patients with COPD with coexisting Lung Diseases and co-morbidities.

COPD represents an important non-communicable health problem that should be diagnosed properly before treatment and to determine a management plan. COPD is a major cause of chronic morbidity and mortality throughout the world; many people suffer from this disease for years and die prematurely from it or its complications. Globally, the COPD burden is projected to increase in the upcoming decades because of continued exposure to risk factors and association of other coexisting illnesses responsible for deterioration of disease<sup>9-13</sup>. Recent study shows that DEF is better than SEF in the management of COPD in LMICs.

Disease, Exacerbation and FEV<sub>1</sub> (DEF) based classification is more appropriate than Symptom, Exacerbation and FEV<sub>1</sub> (SEF) based classification for COPD management<sup>1</sup>

## 1.2 Management strategy of COPD<sup>1</sup>:

According to WHO management Strategy of COPD, an effective COPD management plan includes four fundamental components:

- (1)** assess and monitor disease;
- (2)** reduce risk factors;
- (3)** manage stable COPD;
- (4)** manage exacerbations of COPD.

The goals of effective COPD management are to:

- o Prevent disease progression by controlling Risk factors
- o Relieve symptoms by modern Pharmacological medicine
- o Improve exercise tolerance by Pulmonary Rehabilitation Program
- o Improve health status by controlling coexisting diseases and comorbidities
- o Prevent and treat complications by precise diagnosis of all components of COPD
- o Prevent and treat exacerbations by vaccination and avoiding air pollution, temperature changes and early detection and treating infection
- o Reduce mortality by reducing exacerbations, hospitalizations frail syndrome and by controlling coexisting diseases and comorbidities.

## 2 CHAPTER 2: METHODOLOGY

Global Initiative for Chronic Obstructive Lung Disease (GOLD) guideline has been employed from 2001 to till 2008 in Bangladesh. We made our national prevalence survey BOLD BD study 2007<sup>14</sup>.

In 2008 we developed and published a COPD guideline based on Symptom, Exacerbations and Function (SEF) classification for COPD<sup>10</sup>. However, disease-based classification (DBC) appears to be a better than the SEF. That's

There exists under-reporting of productive cough as most patient swallow their sputum. As such, symptom-based classification might be misleading in the management of COPD

why we worked on DBC and developed a new thought which includes the diagnosis of every component of the 'disease cluster' of COPD and formulate an individualized-tailored management protocol.

### 2.1 Guideline development:

We conducted an extensive search on PubMed using the search term 'COPD', AND/OR 'Chronic Obstructive Pulmonary Disease' with a limit to 'abstracts', 'Clinical Trial' or 'Meta-analysis study type' and COPD related review papers. We have also conducted manual search for NICE guideline, Australian Guideline, GOLD 2020 and SEF Bangladeshi guidelines. The pre-defined selected articles were evaluated by Bangladesh COPD guideline committee. We have used The Appraisal of Guidelines for Research & Evaluation (AGREE) II Instrument for evaluation of various guidelines and review papers. From review of different guidelines and papers our committee made an evidence-based consensus statement which is, supported by various papers and/ or guidelines. Moreover, the guideline is anchored on the background of our relevant published review paper<sup>15</sup>.

## 3 CHAPTER 3: DEFINITION

### 3.1 Definitions in short

**COPD:** Chronic Obstructive Pulmonary Disease (COPD) is a cluster of at least two or more pulmonary diseases that is characterized by persistent **exertional dyspnea and/ or chronic cough with or without expectorations** and airflow limitation, and that is due to airway and/or alveolar abnormalities with or without Pulmonary hypertension usually induced by significant exposure to noxious particles or gases.


**Cluster of disease:** Aggregation of different disease entities is various phenotypes of COPD is defined as ‘cluster of COPD’ in this guideline e.g. ‘Chronic Bronchitis + Emphysema’, ‘Emphysema + Asthma’, ‘Chronic Bronchitis + Emphysema + GERD’ etc.

**Co-morbidity:** The term “comorbidity” has been used to indicate a medical condition existing simultaneously with but independently of another condition.

**Co-existing disease:** Although comorbidity and coexisting diseases terminology is synonymous but, in this guideline, coexisting diseases are defined as the conditions which develop as a consequence during progression and raising degree of severity of COPD.

**Persistent respiratory symptoms:** Exertional dyspnea and/ or chronic cough ± expectorations are defined as persistent respiratory symptoms in this guideline.

Note: by definition pure chronic bronchitis, Pure Pulmonary Emphysema and pure asthma even severe persistent asthma is not included in the definition of COPD, as by definition COPD is a cluster of at least 2 or more diseases. For example.



COPD, in most of the cases is a cluster of, “chronic bronchitis and emphysema”, which is not included in the definition<sup>16</sup>,

Figure 2: Clusters of COPD phenotype



**Emphysema**, by definition is destruction of the gas-exchanging surfaces of the lung (alveoli), is a pathological term that is often (but incorrectly) used clinically and describes only one of several structural abnormalities present in patients with COPD<sup>17</sup>.

**COPD-S** means stable COPD i.e. Usually this group of COPD patient having no coexisting lung disease.

**Unstable COPD:** (COPD – U); COPD U means COPD patient having more than two exacerbations per year without hospitalization or one exacerbation with hospitalization per year

**COPD-C** means COPD patient having at least one coexisting lung disease. Two or more coexisting lung disease may be present in an individual COPD patient

### 3.2 Clinical definition of COPD:

COPD is defined as above, and it's various clusters are described in the figure below:




Figure 3: Non-proportional Phenotypes of COPD  
 (Reproduced by the permission of ATS)

Although non-proportional Venn diagram represents only the component of the cluster of disease however, by proportional diagram we can understand the relative burden of components of the cluster of COPD as shown in figure below.




Figure 4: Proportional Venn Diagram of COPD

(Reproduced by seeking permission) Proportional Venn diagram presenting the different phenotypes within the Wellington Respiratory Survey study population<sup>18</sup>.

**Few facts and findings** of COPD are:

- Despite sputum production being a common symptom, usually it is under-reported by most of the patients<sup>3</sup>.
- The main risk factor for COPD is tobacco smoking but other environmental exposures such as biomass fuel exposure and air pollution may contribute.
- Furthermore, exposure to smoke and/or smoking, host factor influences are also important to develop COPD. These include genetic abnormalities, abnormal lung development and accelerated aging<sup>3</sup>.
- Lung function usually deteriorate after exacerbations and usually never return to previous state after each exacerbation in COPD as shown in the figure below:




Figure 5: Disease trajectory of COPD

A stylized model depicting the end-of-life trajectory for a proportion of patients with COPD<sup>19</sup>

COPD may be associated with coexisting pulmonary diseases and significant other comorbid chronic diseases, which increase its morbidity and mortality<sup>2,15</sup>.

After commencement of COPD, the lung function progressively worsens with increasing severity and this is determined by the inducing factors i.e. smoke, host factors, abnormal lung development, recurrent infection, allergic inflammation and hypoxic spell during sleep and during exacerbations.

Significant coexisting pulmonary diseases and comorbidities may have an impact on morbidity and mortality<sup>2</sup>. There may be significant lung pathology (e.g. emphysema) in the absence of airflow limitation that needs further evaluation as simple emphysema is not defined as COPD.

The chronic and non-reversible or partially reversible airflow limitation that is characteristic of COPD is caused by a mixture of small airways disease and parenchymal destruction (emphysema), the relative contributions of which vary from person to person<sup>20</sup>.

These changes do not always occur together at a time but evolve at different rates over time. Chronic inflammation causes structural changes, narrowing of the small airways and destruction of the lung parenchyma that leads to the loss of alveolar attachments to the small airways and decreases lung elastic recoil. In turn, these changes diminish the ability of the airways to remain open during expiration. A loss of small airways may also contribute to airflow limitation

and mucociliary dysfunction is a characteristic feature of the disease<sup>21</sup>. Airflow limitation is usually measured by spirometry as this is the most widely available and reproducible test of lung function<sup>22</sup>.

Exertional shortness of breath, barrel shaped chest, obliteration of 'area of cardiac dullness', low down diaphragm and silent chest are the diagnostic features of emphysema component of COPD.

Chronic bronchitis<sup>23</sup>, by definition is presence of cough and sputum production for at least 3 months in each of two consecutive years, still remains a clinically and epidemiologically useful term with spirometrically normal lung function. This definition is usually not used during diagnosis of COPD.

After widespread small airway obstruction, changes in spirometry measurement will show  $FEV_1/FVC \% < 0.7$  or lower limit normal (LLN) and  $FEV_1$  80% or less<sup>24</sup>. These changes are defined as Chronic Obstructive Bronchitis, an important component of early COPD.

In COPD, presence of chronic cough and sputum production for more than 2 years is a useful diagnostic point of chronic bronchitis component of COPD. Persistent respiratory symptoms also exist in individuals showing normal spirometry<sup>25</sup> and a significant number of smokers without airflow limitation have structural evidence of lung disease manifested by the varying presence of emphysema, airway wall thickening and gas trapping. They are by definition not COPD.

### **3.2.1 Pre-COPD or Early COPD<sup>26</sup>:**

These are group of patients who have history of significant exposure to smoke and/or smoking along with some of the early symptoms of COPD include coughing with or without expectorations, excess mucus, shortness of breath, and tiredness or structural changes clearly like emphysema like hyperinflated lung but spirometry showing  $FEV_1/FVC \% > 0.7$  or LLN and  $FEV_1$  80% or more. Identifying these patients is imperative to prevent progression of disease.

## 4 CHAPTER 4: DEVELOPMENT AND PROGRESSION of COPD

Cigarette smoking is the most well studied COPD risk factor for commencement of disease. But it is not the only risk factor and there is consistent evidence from epidemiologic studies that non-smokers may also develop chronic airflow limitation<sup>27</sup>. Much of the evidence concerning risk factors for COPD comes from cross-sectional epidemiological studies that identify associations rather than causal relationships. Nevertheless, compared to smokers with COPD, nonsmokers with chronic airflow limitation have fewer symptoms, milder disease and lower burden of systemic inflammation<sup>28</sup>. Interestingly, non-smokers with chronic airflow limitation do not appear to have an increased risk of lung cancer, or cardiovascular comorbidities, compared to those with chronic airflow limitation from smoking. That's why tobacco cessation is warranted in smokers with COPD<sup>28</sup>.

However, there is evidence that some non-smoker COPD have an increased risk of pneumonia and mortality from respiratory failure<sup>29</sup>. Biomass fuels (coal, wood, dung, crop waste etc.) exposure is important risk factors in rural and suburban areas of Bangladesh<sup>30</sup>. Occupational dust exposure might be responsible for 20 to 30% of COPD<sup>31</sup>. A summary of the risks of COPD associated with biological or mineral dusts, gases, fumes / vapors, diesel exhaust, chemical gas / fumes and various other occupational exposures appears in Figure 4<sup>32</sup>




Figure 6: Risk of occupational exposure for COPD

Risk of occupational exposure for COPD adapted from Australian and New Zealand Guidelines 2019

Biological dust exposures include substances of microbial, plant or animal origin such as bacteria, fungi, allergens, endotoxins, peptidoglycans, glucans, pollens, and plant fibers<sup>33</sup>.

This occupational risk factors study clearly demonstrated that biological dust, Diesel exhaust, mineral dust, irritant gases and vapors, Dusts (PM 10 and PM 2.5) and chemical gas and fumes have significant association for development of COPD<sup>33-37</sup>.

The development of COPD is most likely a series of complex interactions between noxious airborne agents together with a genetic predisposition<sup>38</sup>. Exposure to cigarette smoke and air pollution as well exposure to infectious agents are commonly associated with the development of COPD<sup>39</sup>. Because COPD does not develop in all smokers and it is interesting that even for heavy smokers, fewer than 50% develop COPD during their lifetime. Genetic predisposition such as protease-antiprotease imbalance is a plausible explanation for the development of airways inflammation and airflow obstruction<sup>40</sup>.

## 5 CHAPTER 5: OVERVIEW

Chronic Obstructive Pulmonary Disease (COPD) is actually a cluster of diseases associated with progressive airflow limitation which is not fully reversible. Chronic inflammatory process throughout the airways, parenchyma & pulmonary vasculature resulting in many structural and functional changes in the lung. COPD is a burden for both developed and developing countries. COPD management is a great challenge even after implementing COPD guidelines properly<sup>41</sup>.

The term “comorbidity” has been used to indicate a medical condition existing simultaneously with but independently of another condition<sup>42</sup>. Recently, however, use of the term comorbidity has broadened to suggest a causal relationship between two disease states. Certainly, in case of chronic obstructive pulmonary disease (COPD), assessment of all other components of disease is very important from both an academic and a clinical point of view. It is likely that infection has a larger role than currently recognized in the pathogenesis of COPD, and the relationship between the two can be viewed as a complicated comorbid one, which may affect both the direction and course of each problem<sup>43</sup>.




Figure 7<sup>43</sup>: Two distinct infection cycles in COPD

A new coexisting disease<sup>44</sup>-based classification is formulated which is directed more towards management of COPD. It's based on a self-made theme quotation “If Arterial Blood Gas (ABG) deteriorates, stable COPD Patient may Turn Out (PTO) to unstable COPD”.

Although comorbidity and coexisting diseases terminology is synonymous but, in this guideline, coexisting diseases to describe: few definite conditions which develop during progression and stage changes of COPD<sup>45</sup>. These coexisting diseases may be developed as a sequel or consequence of progression of COPD which leads to more frequent exacerbations and changes from stable status of COPD to COPD with coexisting diseases and then gradually towards unstable COPD.

It is now clear that stable COPD patient due to continuous exposure to smoke or smoking with recurrent infection, bronchospasm, exposure to air pollutant and temperature changes with genetic and hormonal susceptibilities changes gradually progresses to unstable COPD over the years. That's why our theme quotation is very simple to recapitalized with the acronym 'ABG' and 'PTO'. It is now clear that some internal and external factors are responsible for turning out stable COPD patient's to unstable COPD<sup>9</sup>. If we can control six important factors which are recognized as triggers for exacerbation of COPD, exacerbation may be curtailed and may be prevented. Out of six factors four are internal factors and two are external factors.

We can use mnemonic 'ABG' and 'PTO' for the risk factors for COPD. That's why we used the theme quotation. (Table 1)

| RISK FATOR                              |   |   |                                |
|---|---|---|--------------------------------|
| Asthma                                  | A | P | Pollution                      |
| Bronchiectasis                          | B | T | Temperature changes            |
| Gastro-Esophageal Reflux Disease (GERD) | G | O | Obstructive Sleep Apnea (OSA). |

Table 1: Mnemonic meaning of 'ABG' & 'PTO'

Here actually 'A' means Asthma, 'B' means Bronchiectasis, G means Gastro-Esophageal Reflux Disease (GERD), 'P' means air Pollution, 'T' means Temperature changes and 'O' means Obstructive Sleep Apnea (OSA).

Precisely two steps for commencement and progression of COPD: initiation of inflammation from interaction with smoke/ smoking plus genetic factors along with recurrent infection – both are responsible for development and progression of COPD<sup>38,41,43,46</sup>. Asthma, GERD, air pollution, temperature changes and OSA -all developed during the progress of COPD are responsible to perpetuate airways and lung parenchymal inflammation. Recurrent infection in patients of COPD is the main cause that initiates and maintain neutrophilic inflammation, thick secretion, peripheral airways obstruction. This persistent



neutrophilic inflammation ultimately causes necrotizing inflammation and destruction of the airways smooth muscle that perpetuate to much more peripheral bronchiectasis.

## 6 CHAPTER 6: DEF CLASSIFICATION AND STAGING

DEF Classification of COPD is as follows<sup>15</sup>:

1. Stable COPD: (COPD- S) i.e. Indolent COPD, having no coexisting lung disease with having one or less exacerbation per year and virtually no hospitalization for COPD

2. COPD with coexisting diseases: COPD-C

- COPD with Asthma: COPD-A
- COPD and Bronchiectasis: COPD-B
- COPD and GERD: COPD-G
- COPD and Obstructive Sleep Apnea (OSA): COPD-O

3. Unstable COPD: (COPD – U); COPD U means COPD patient having more than two exacerbations per year without hospitalization or one exacerbation with hospitalization per year

COPD-S, COPD-C and COPD-U:

Still there is no internationally accepted definition of Stable or Unstable COPD. We developed a working definition for Stable or Unstable COPD for patient management. It is defined as per follows:

- COPD-S means stable COPD i.e. Usually this group of COPD patient having no coexisting lung disease. That means COPD patient having no feature of Asthma, Bronchiectasis, OSA or GERD. Progression of disease in Stable COPD patient depends mainly on temperature change, air pollution and/ or Smoking effect. That's why nature of progression would be slow with infrequent exacerbation, i.e. less than two exacerbations per year and virtually no hospitalization in a year for COPD.

- COPD C means COPD patient having at least one coexisting lung disease. Two or more coexisting lung disease may be present in an individual COPD patient. For example, asthma and bronchiectasis may coexist simultaneously, which may be designated as COPD C: AB subtypes or subcategories. Thereby COPD C: subtypes may be presented as single (A/B/ G/O) or as different subtype combinations. That's why in COPD-C group, in addition to 4 single sub-categories, COPD AB, or AG or AO or BG or BO or GO or ABG or ABO or AGO or BGO or ABGO – total 11 possible combinations of multiple coexisting illness may be present in an individual. All the COPD C patients are prone to repeated exacerbations and vulnerable to become COPD U.

- COPD U patients are group of patients who experience repeated exacerbations. Any one of 15 subtypes of COPD C group may turn to COPD U if they develop frequent exacerbations or hospitalization in a year.

## 6.1 Distinct pathophysiological features:

Four distinct pathophysiological features are crucial for development and perpetuation of unstable COPD: i.e. recurrent exacerbations and hospitalization

1. Recurrent or persistent Eosinophilic Inflammation<sup>47,48</sup>: Features like Asthma.
2. Recurrent infection and neutrophilic inflammation<sup>11</sup>: Features like Bronchiectasis, para-septal Bullae are likely to develop.
3. Acid/Bile aspiration and micro-aspiration of food particles, throat content with or without micro-organism with neutrophilic inflammation<sup>49</sup>: Features like recurrent right and left basal pulmonary inflammation, distortion of small airways and to some extent pulmonary fibrosis is likely to develop.
4. Hypoxic spell during sleep and during exacerbations may lead to development of pulmonary hypertension<sup>50</sup>




Figure 8: A Journey of Stable to Unstable COPD

For precise diagnosis and sub-categorization (single or combinations) of COPD C, we need some investigations i.e. a good quality Chest X-ray, HRCT scan of chest (if possible), CBC, and total circulating eosinophil count, sputum cytology with differential count, Blood total IgE level and FeNO<sup>51</sup> [preferably at least 2 weeks not getting oral Corticosteroids (OCS) or inhaled Corticosteroids (ICS). If patient is getting OCS and/or ICS then at least 2-3 times FeNO should be done in 1 to 3 months interval to find out rising trends of FeNO in COPD.




Figure 9<sup>52</sup> Coexisting diseases and comorbidities in COPD.

## 6.2 Cardio-Pulmonary Exercise Test (CPET)

It is important for the management plan of COPD particularly for rehabilitation program of COPD<sup>53</sup>. As it needs a costly Ergometry machine, we developed Modified CPET (M-CPET) test, that means a combination of Spirometry with reversibility test, Incremental Shuttle walk test<sup>54</sup>, pre-exercise ECG and 6-minute cycling with Tele-cardiac monitoring<sup>55</sup>.

Modified CPET (M-CPET) can't check anaerobic threshold (AT) and  $VO_2$  Max. But this can measure lung function with reversibility status effectively, exercise capacity before and after rehabilitation program and find out evidence of cardiac ischemia and different types of arrhythmias during exercise. This test will measure Level of increment instead of  $VO_2$  Max.

Spirometry is important for staging of COPD<sup>56</sup> and Modified CPET starts with spirometry with reversibility testing, this test is also important to diagnose and classify stage of COPD as well as simultaneously diagnosis of COPD– A sub-group. In COPD-A reversibility test may be partially positive, i.e. either improvement of  $FEV_1$  at least 200ml alone or at least 12% improvement of  $FEV_1$ .

## 6.3 SEVERITY OF COPD:

| <b>Classification of airflow limitation severity in COPD</b><br>(Based on post-bronchodilator FEV <sub>1</sub> , in patients with FEV <sub>1</sub> /FVC Ratio <0.7) |             |  |
|---|-------------|--|
| STAGE 1   | MILD        | FEV <sub>1</sub> ≥ 80% PREDICTED       |
| STAGE 2   | MODERATE    | 50% ≤ FEV <sub>1</sub> < 80% PREDICTED |
| STAGE 3   | SEVERE      | 30% ≤ FEV <sub>1</sub> < 50% PREDICTED |
| STAGE 4   | VERY SEVERE | FEV <sub>1</sub> < 30% PREDICTED       |

Table 2: Severity of COPD on the basis of FEV1 GOLD 20203.

In some COPD cases Spirometry may not meet criteria FEV<sub>1</sub>/FVC ratio <0.7, due to presence of other coexisted diseases or comorbidities, obesity, presence of multiple bullae, chest wall muscle or diaphragmatic muscle weakness, electrolyte imbalance etc<sup>57</sup>.

By definition in COPD patient, airflow limitation is not fully reversible, after administration of bronchodilator medication, the ratio of FEV<sub>1</sub> to forced vital capacity (FVC) is <70% even the FEV<sub>1</sub> is ≥80% of the predicted value. The ratio of FEV<sub>1</sub> to vital capacity (VC) is a sensitive indicator for mild COPD. FEV<sub>1</sub>/FEV<sub>6</sub> has a high level of agreement with FEV<sub>1</sub>/FVC on both the fixed ratio and Lower Limit of Normal (LLN) criteria for the diagnosis of COPD. There is controversy regarding the optimal cut-off to define airflow limitation (FEV<sub>1</sub>/FVC less than 0.7 versus lower limit of normal). There is evidence that the fixed ratio can lead to over diagnosis of COPD in older populations, under diagnosis in younger people<sup>58-60</sup> and may lead to gender imbalances as women have higher FEV<sub>1</sub>/FVC ratio than their male counterparts<sup>61</sup>. A systematic review of 11 studies which examined the relationship of each criterion with clinical outcomes found both were related to clinical outcomes and concluded that on current evidence one could not be preferred over the other. The LLN appeared to be a better criterion in older patients with less severe airflow limitation<sup>57</sup>; however, a study shows that the fixed cut-off of 0.7 identified more people with CT diagnosed emphysema<sup>62,63</sup>.

## 7 CHAPTER 7: DIAGNOSIS

Diagnosis part has Three Components:

- Diagnosis of COPD
- Diagnosis of Coexisting Diseases
- Diagnosis of Comorbidities

### 7.1 Summary notes:

COPD should be considered in any patient who has dyspnea, chronic cough or sputum production, a history of recurrent lower respiratory tract infections and/or a history of exposure to risk factors for the disease.

Spirometry is required to make the diagnosis; the presence of a postbronchodilator  $FEV_1/FVC < 0.70$  confirms the presence of persistent airflow limitation although various contributing factors may limit the presence of a post-bronchodilator  $FEV_1/FVC\% < 0.70$ . Under 50 years age we should practice lower limit normal (LLN)  $FEV_1/FVC\%$ . Always  $FEV_1$  is important to understand stage of COPD and consider limitation of  $FEV_1/FVC$  ratio  $<0.70$  under 50 years of age.

After diagnosis of COPD, diagnosis of coexisting illnesses would be established for sub-classification of COPD.

The goals of COPD assessment are to determine the level of airflow limitation, the impact of disease on the patient's health, and the risk of future events (such as exacerbations, hospital admissions, or death), in order to guide therapy.

Concomitant chronic diseases occur frequently in COPD patients, including cardiovascular disease, skeletal muscle dysfunction, metabolic syndrome, osteoporosis, depression, anxiety, and lung cancer.

Coexisting diseases and comorbidities should be actively sought and treated appropriately when present as they can influence morbidity, hospitalization, and mortality independently.





Figure 10: Algorithm of diagnosis of COPD and Coexisting disease





Figure 11: Algorithm for Management of COPD



CAT score or / mMRC Score: any one score can be used during follow up of patient and to understand level of symptom control and pre and post Pulmonary Rehab assessment.

Figure 12: CAT SCORE<sup>64</sup>

| CAT™ ASSESSMENT  |  |  |
|--|--|--|
| For each item below, place a mark <input type="radio"/> in the box that best describes you currently. Be sure to only select one response for each question. |  |  |
| EXAMPLE: I am very happy   | <input type="radio"/> 0 <input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | I am very sad  |
| I never cough  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | I cough all the time   |
| I have no phlegm (mucus) in my chest at all  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | My chest is completely full of phlegm (mucus)                          |
| My chest does not feel tight at all  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | My chest feels very tight  |
| When I walk up a hill or one flight of stairs I am not breathless  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | When I walk up a hill or one flight of stairs I am very breathless     |
| I am not limited doing any activities at home  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | I am very limited doing activities at home                             |
| I am confident leaving my home despite my lung condition   | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | I am not at all confident leaving my home because of my lung condition |
| I sleep soundly  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | I don't sleep soundly because of my lung condition                     |
| I have lots of energy  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5            | I have no energy at all  |
|  |  | TOTAL SCORE: <input type="text"/>                                      |

Figure 13: CAT SCORE (Bangla Version)

| CAT SCORE (BANGLA VERSION)   |   |   |
|--|---|---|
| আপনার নাম :  | আজকের তারিখ :   |   |
| আপনার সিঁগপিডি অসুখটি কোন পর্যায়ে আছে? সিঁগপিডি অসুখটির অবস্থা যাচাই করুন।  |   |   |
| নিম্নোক্ত প্রশ্নাবলী আপনাকে ও আপনার স্বাস্থ্য সেবা প্রদানকারীকে আপনার ভাল থাকা ও প্রাত্যহিক জীবনে সিঁগপিডি (ক্রনিক অবস্ট্রাক্টিভ পালমোনারী ডিজিজ) অসুখটির প্রভাব পরিমাপে সহায়তা করবে। আপনার উত্তর ও আপনার প্রাণ ফোর আপনাকে ও আপনার স্বাস্থ্যসেবা প্রদানকারীকে আপনার চিকিৎসার উন্নয়নে ও তা থেকে সর্বোচ্চ উপকার কিভাবে পাওয়া যাবে তা নির্ণয়ে সহায়তা করবে। |   |   |
| নীচের প্রতিটি আইটেম এর জন্য, যে কষ্টটি আপনার বর্তমান অবস্থা বর্ণনা করবে তাতে টিক (✓) চিহ্ন দিন। প্রতিটি প্রশ্নের জন্য একটি মাত্র উত্তর দান নির্ধিত করুন।   |   |   |
| উদাহরণঃ আমি খুবই সুখী  | <input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5              | আমি খুবই বিষ্ময়।                                       |
| আমার কাশি হয়না  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | আমার সর্বদাই কাশি হয়                                   |
| আমার বুকে শ্বেদা জমে নেই   | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | আমার বুকে শ্বেদায় পরিপূর্ণ                             |
| আমি বুকে মোটেও চাপ অনুভব করিনা   | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | আমি বুকে প্রবল চাপ অনুভব করি                            |
| যখন ঢাল বা সিঁড়ি বেয়ে ওপরে উঠি তখন মোটেও শ্বাসকষ্ট হয়না   | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | ঢাল বা সিঁড়ি বেয়ে উপরে ওঠার সময় প্রবল শ্বাসকষ্ট হয়  |
| আমি সব গৃহকর্মই করতে পারি  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | আমি কোন গৃহকর্ম করতে পারিনা                             |
| আমার শ্বাসতন্ত্রের সমস্যা সত্ত্বেও আত্মবিশ্বাস নিয়ে বাড়ির বাইরে যাই  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | শ্বাসতন্ত্রের সমস্যার জন্য আমি বাড়ির বাইরে যেতে পারিনা |
| আমার সুনিদ্রা হয়  | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | আমার মোটেও সুনিদ্রা হয় না                              |
| আমার প্রচুর প্রাণশক্তি   | <input type="radio"/> 0 <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 | আমার মোটেও শক্তি নেই                                    |

সর্বমোট কোরঃ

Table 3: Modified Medical Research Council (mMRC) Scale Grading the severity of breathlessness during daily activities<sup>65</sup>

| Grade Description of Breathlessness  |
|--|
| Grade 0: I only get breathless with strenuous exercise   |
| Grade 1: I get short of breath when hurrying on level ground or walking up a slight hill   |
| Grade 2: On level ground, I walk slower than people of the same age because of breathlessness, or I have to stop for 2 breath when walking at my own pace on the level |
| Grade 3: I stop for breath after walking about 100 metres or after a few minutes on level ground   |
| Grade 4: I am too breathless to leave the house or I am breathless when dressing or undressing   |

Table 4: Modified Medical Research council scale in Bangla

| Mdified MRC Scale: (MRC Score) |   |
|--------------------------------|---|
| Score                          | Scale   |
| 0                              | প্রচণ্ড পরিশ্রম ছাড়া শ্বাসকষ্ট হয়না।  |
| 1                              | সমতলে দ্রুত হাঁটলে অথবা সামান্য উঁচুর দিকে হাঁটলে শ্বাসকষ্ট হয়।                          |
| 2                              | সমবয়সী লোকদের তুলনায় আন্তে হাঁটতে হয়।  |
| 3                              | শ্বাস নেয়ার জন্য সমতলে ১০০ গজ হাঁটার পরে অথবা কয়েক মিনিট পরেই বিশ্রাম নিতে হয়।         |
| 4                              | এত শ্বাসকষ্ট যে বাড়ি থেকে একা বের হতে সাহস হয়না অথবা জামা কাপড় পরতে গেলে হাঁপিয়ে উঠি। |

## 8 CHAPTER 8: INVESTIGATIONS<sup>63</sup>

### 8.1 CBC:

Complete blood count is important to understand

- Polycythemia (More in Ch bronchitis predominant COPD)
- Anemia (More in Emphysema predominant COPD)
- Higher count with High polymorph may indicate RTI
- Eosinophil  $\geq 300$  per  $\mu\text{L}$  indicating, may need ICS therapy
- Low MCV, MCH with anemia indicating Iron deficiency anaemia.

### 8.2 Chest X-Ray

A plain Postero-Anterior and Lateral Chest x-ray helps to exclude other conditions such as lung cancer. Limitation of CXR are non-specificity of lung hyperinflation and identifying nodules  $< 1$  cm. But Chest X-Ray is important to identify other coexisting diseases e.g.

- Bronchiectasis, Lung Cancer, Lung Fibrosis etc.
- Pneumonia or Respiratory tract infection old PTB with fibrosis etc.
- Pneumothorax, Bulla
- Cardiomegaly or Tubular heart
- Emphysematous changes

### 8.3 Sputum examination

Sputum culture is recommended when an infection is not responding to antibiotic therapy or when a resistant organism is suspected. Sputum examination is recommended to understand

- Sputum Eosinophil:  $\geq 3$  or more per HPF; indication ICS therapy in COPD
- Sputum Neutrophil: plenty in Bronchiectasis, Lung infection with COPD and need of Pulse Azithromycin therapy
- Sputum Gene X-pert and AFB for Typical and/or atypical Mycobacterial Infection
- Sputum culture during exacerbation to identify drug-resistant organism.

### 8.4 Spirometry

Spirometry is the most reproducible and objective measurement of airflow limitation. It is a noninvasive and readily available test.




Figure 14: Comparison of different flow-volume curves

The spirometry is very much dependent on patient effort and cooperation, and there are important quality criteria that should be met in conducting spirometry<sup>60</sup>.

#### Indications for Spirometry:

- Breathlessness that seems to be inappropriate;
- Chronic (daily more than 3 weeks) cough or intermittent,
- Unusual cough;
- Frequent or unusual sputum production
- Relapsing acute infective bronchitis; and
- History of risk factors such as exposure for COPD.
- Strong family history of COPD or Asthma

### 8.5 Exercise testing

Exercise testing is carried out to find out three questions:

1. Is exercise capacity normal?
2. What is the cause of abnormal exercise capacity?
3. Is exercise training safe to perform?




Figure 15: Causes of breathlessness

Three principal causes of exercise intolerance are mainly identified, ventilatory limitation, Cardiac limitation, and Muscular limitation.

Cardio-Pulmonary Exercise Tasting (CPET) is the gold-standard, however, many facilities have no such expensive cycle ergometer for the test. We developed a 'Modified Cardio-Pulmonary Exercise Tasting (CPET)' for our Bangladeshi Patients:

### 8.5.1 Components of Modified CPET:

- Spirometry with Reversibility test
- Resting ECG
- Incremental shuttle walk test or 6-minute walk Test
- Tele- Cardiac Monitoring (TCM) before, during and after Exercise

We do Modified CPET test to find out-

- Baseline (Before Rehab) and after Rehab Lung Function (Pre and post bronchodilator  $FEV_1$ , FVC and  $FEV_1/FVC$  ratio)
- Patient Endurance before and after Rehab by Incremental shuttle walk test or 6-minute walk Test

- Cardiac arrhythmias and ischemia before, during and after exercise

Why for Bangladeshi patient, develop this test protocol instead of Standard Ergometry?

- Ergometry is a costly machine (usually not possible to purchase by most of physician and it is available only in few referral center of Bangladesh)
- To Understand clearly patients pre-Rehab and post-Rehab Endurance (Stamina)
- To understand Lung Function changes before and after Rehabilitation
- To prevent patients' Cardiac Catastrophe by early diagnosis of Ventricular Extra Systole with the help of Tele-Cardiac Monitoring
- Incremental shuttle walk test (ISWT) or 6 Minute Walk Test (6MWT) can be used by primary care physician

Therefore, Followings are the indications of M-CPET

- Pre-Rehab assessment of patient
- To understand and to explain the real endurance (stamina) and CardioPulmonary Function of patient
- Post -Rehab assessment of patient
- To assess patient before Cardio-Pulmonary Surgery

**Note: Physician may use 6 Minute walk test<sup>66</sup> or Incremental Shuttle walk test<sup>67-70</sup> and other components of M-CPET separately for patient assessment**

## 8.6 Fractional exhaled Nitric Oxide (FeNO)<sup>51,71</sup>

FeNO should be tested to all COPD patients to prescribe Inhaled Corticosteroids (ICS) appropriately

- Differentiation of types of airway inflammation is very important in the management of chronic respiratory illnesses like COPD having with or without asthma.
- Presence of eosinophilic inflammation means Inhaled Corticosteroids (ICS) should be used in the management of asthma and COPD.
- Presence of Non-eosinophilic Inflammation means only bronchodilator is enough to control symptom; i.e. ICS has little role to control inflammation of COPD.

### What is FeNO breathing test?

This is a simple, quantitative, noninvasive way to measure eosinophilic airways inflammation i.e. FeNO is a surrogate biomarker of eosinophilic inflammation.

### Advantages of FeNO:

1. Detects eosinophilic airway inflammation- justifies use of Inhaled Corticosteroids (ICS)
2. It can determine likelihood of corticosteroid responsiveness
3. Can be used as a monitoring tool in ICS treated patients

Interpretation of FeNO (ATS Recommendation)<sup>72</sup> Levels under 20 parts per billion (PPB) in children and under 25 parts per billion (PPB) in adults are considered normal.

### Japanese Recommendation<sup>73,74</sup>:

1. FeNO NO less than 22 ppb– Minimum eosinophilic inflammation and responsiveness to inhaled corticosteroids is less likely
2. FeNO 22 ppb or more- indicate eosinophilic inflammation and likely to responsiveness to Inhaled corticosteroids

FeNO- in monitoring: ATS Recommendation<sup>72</sup>:

1. Significant increase in FeNO: > 20% for values over 50 ppb or more than 10 ppb for values < 50 ppb from one visit to the next
2. Significant response to anti-inflammatory therapy: Fall at least 20% of FeNO for Values more than 50 ppb or fall at least 10 ppb for values lower than 50 ppb from one visit to the next

For this reason, one can advise to do FeNO at random, if result is inconclusive or very low, should stop ICS. If patient is symptomatic or after 2 months recheck FeNO to find out relapse of eosinophilic inflammation. If FeNO level increases, indicating need to use Inhaled Corticosteroids (ICS).

## 8.7 Saturation of pulsatile O<sub>2</sub> (SpO<sub>2</sub>)<sup>75-77</sup>

Oximeters typically have an accuracy of  $\pm 2\%$ , which is satisfactory for routine clinical practice. They are more useful for monitoring trends than in single

measurement. Oximetry does not provide any information about carbon dioxide status and is inaccurate in the presence of poor peripheral circulation (e.g., cold extremities, cardiac failure) and when SpO<sub>2</sub> readings are consistently below 90%.

Table 5<sup>78</sup>: Calculation for PaO<sub>2</sub> assessment as

| SpO <sub>2</sub> | Calculation for PaO <sub>2</sub>   | Resultant PaO <sub>2</sub> range |
|------------------|--|----------------------------------|
| 100%-90%         | Decrease PaO <sub>2</sub> by 4 mmHg for every single percent reduction in SpO <sub>2</sub>   | 100-60 mmHg                      |
| 90%-80%          | Decrease PaO <sub>2</sub> by 1.5 mmHg for every single percent reduction in SpO <sub>2</sub> | 60-45 mmHg                       |
| <80%             | Decide SpO <sub>2</sub> by 2 to reach to a PaO <sub>2</sub> level                            | ≤ 40 mmHg                        |

This formula, which is not an exact mathematically proven entity, can be of extreme help to ICU doctors and Respiratory practitioners. For example, if the monitor is showing a SpO<sub>2</sub> of 70%, we can almost consider a value of PaO<sub>2</sub> to be around 35 mmHg and take appropriate measures for the patient.

This hypothesis can have some pitfalls, for example, this calculation will not be effective in cyanide poisoning and certain hemoglobinopathies, but still, a fair and working assessment may be drawn from this calculation.




Figure 16<sup>78</sup>: Oxyhemoglobin dissociation curve

## 8.8 Arterial blood gas measurement<sup>76-80</sup>

Arterial blood gas analysis should be considered in all patients

- with severe COPD
- when considered for domiciliary oxygen therapy (e.g., whose FEV<sub>1</sub> is <40%)

predicted or  $<1$  L, whose oxygen saturation as measured by pulse oximetry [ $SpO_2$ ] is  $<92\%$ )

- those with pulmonary hypertension
- those with breathlessness out of proportion to their clinical status
- Respiratory failure is defined as a  $PaO_2 < 60$  mmHg (8 kPa) with or without  $PaCO_2 > 50$  mmHg (6.7 kPa). The latter is termed ‘ventilatory failure or hypercapnic respiratory failure’ and is accompanied by either compensated (chronic) or uncompensated (acute) acidosis. Acute respiratory acidosis indicates a need for assisted ventilation.

### 8.9 Biochemistry

- High IgE level may be used in COPD A for Omalizumab therapy<sup>81</sup>
- Hyperthyroidism and acidosis are associated with breathlessness<sup>82,83</sup>.
- Hyperventilation states are associated with respiratory alkalosis<sup>84</sup>.
- Hypothyroidism aggravates obstructive sleep apnea<sup>85</sup>.
- COPD with an association between thrombocytosis (400/ $\mu$ L on admission) and mortality<sup>86</sup>. Thrombocytosis (after controlling for confounders) was associated with an increased 1-year all-cause mortality and an increased in hospital mortality (OR 1.53 (95% CI 1.03 to 2.29,  $p=0.030$ ) and OR 2.37 (95% CI 1.29 to 4.34,  $p=0.005$ ) respectively<sup>87</sup>.
- Severe homozygous (ZZ) alpha<sup>1</sup> antitrypsin deficiency (alpha-AT) has been estimated at between 1/4,348 and 1/5,139 in European populations<sup>88</sup>. No such data available in Bangladeshi population, but 75 to 85% of such individuals will develop emphysema<sup>89</sup>. Genetically pure emphysema without Chronic Bronchitis or asthma should be regarded as genetic disease. It is not COPD.
- Tobacco smoking is still the most important risk factor for COPD<sup>90</sup> even in this group of alpha<sup>1</sup> antitrypsin deficiency. Targeted screening suggests between 1.0 to 4.5% of patients with COPD have underlying severe alpha<sup>1</sup> AT deficiency<sup>91</sup>. The index of suspicion should be high in younger patients with predominantly basal emphysema and a family history of COPD.
- The diagnosis can be made by measuring serum levels of alpha<sup>1</sup> antitrypsin (if available) and if reduced, genotyping should be performed.

### 8.10 E.C.G and Echo<sup>83</sup> ?

Cardiovascular disease is common in patients with chronic obstructive pulmonary disease but is often under-recognized.

- Electrocardiography (ECG) may be useful to alert the clinician to its presence in COPD. In a retrospective Dutch study of patients entering pulmonary rehabilitation, ischemic changes were present on ECG in 21% of all patients and in 14% of those without reported cardiovascular comorbidity.
- Electrocardiography is also indicated to confirm arrhythmias suspected on clinical grounds. Multifocal atrial tachycardia is a rare arrhythmia (prevalence  $< 0.32\%$  of hospitalised patients) but over half the cases reported in the literature had underlying COPD.



- Atrial fibrillation commonly develops when pulmonary artery pressure rises, leading to increased right atrial pressure.

### **Usefulness of Echocardiography in COPD:**

- Pulmonary Hypertension • Chronic Cor-Pulmonale
- Cause of breathlessness is out of proportion to the degree of respiratory impairment • Ischemic heart disease • Embolus in atrium
- Left heart failure. Patients with COPD may have poor quality images on transthoracic examination and transesophageal echocardiography may be more helpful.

Patients with COPD are prone to other conditions associated with cigarette smoking, including accelerated cardiovascular, cerebrovascular and peripheral vascular disease, and oropharyngeal, laryngeal and lung carcinoma. Conversely, there is a high prevalence of COPD among patients with ischemic heart disease, peripheral vascular disease and cerebrovascular disease and smoking-related. These patients should be screened for symptoms of COPD, and spirometry should be performed after diagnosis of smoking related any disorder.

### **8.11 High resolution computed tomography<sup>92-94</sup>**

High resolution computed tomography (HRCT) scanning gives precise images of the lung parenchyma and mediastinal structures.

- The presence of emphysema and the size and number of bullae can be determined. This is necessary if bullectomy or lung reduction surgery is being considered.
- HRCT is also appropriate for detecting bronchiectasis. Vertical reconstructions can provide a virtual bronchogram.
- Helical computed tomography (CT) scans with intravenous contrast should be used in other circumstances, such as for investigating and staging Lung Cancer or diagnosis of Pulmonary Embolism etc.

### **8.12 Sleep studies<sup>95,96</sup>**

It is recommended for COPD patients with the following conditions:

- Suspected of having a coexisting sleep disorder e.g. Obstructive Sleep Apnea (OSA)
- With hypercapnia • Pulmonary hypertension in the absence of daytime hypoxemia
- To diagnose Hypoxic spells during sleep • Right heart failure
- Polycythemia
- Continuous overnight oximetry may be used to assess a need for overnight domiciliary oxygen therapy
- Patients receiving long-term domiciliary oxygen therapy during follow up to assess whether hypoxemia has been adequately corrected.

**9.1 COPD with Asthma: COPD-A**

It is well established that about 40% to 56% COPD patients have asthma along with COPD<sup>9,61</sup>. Typically, asthma is characterized by inflammation predominantly involving eosinophils, whereas COPD is characterized by neutrophilic inflammation<sup>97</sup>. Diagnosis of Asthma with COPD is possible with the help of clinical findings and with the help of some investigations (Table 6) Fractional exhaled Nitric Oxide FeNO and blood eosinophil count have been considered as biomarkers of local and systemic eosinophilic inflammation, which got increased in patients with asthma<sup>98</sup>. Total serum IgE and antigen specific IgE levels are also found elevated in those with allergic asthma<sup>99</sup>.

A prospective clinical observational study showed nearly one third of COPD patients had sputum eosinophilia and the number of eosinophils was significantly correlated to the level of exhaled nitric oxide<sup>100</sup>. Fractional exhaled Nitric Oxide (FeNO) has been described as a marker of asthmatic airway inflammation<sup>101</sup>.

In some COPD patients with asthma (i.e, COPD-A) FeNO level may be normal due to effect of smoking or due to continuous use of inhaled corticosteroid<sup>102</sup> But interestingly some patients have persistent elevation of FeNO levels despite treatment with high dose of corticosteroids<sup>71,103</sup> In this protocol, after diagnosis of COPD, we advise to do FeNO in every cases. If FeNO level is found 20 PPB or more we sub-classify the case as 'COPD-A' and it's a strong indication to prescribe inhaled corticosteroids (ICS) whatever the stage of COPD, from stage 2 to 4. If FeNO level is less than 20 PPB, it is advised not to prescribe inhaled corticosteroids (ICS). It indicates use of only long acting dual bronchodilators and it may be adequate for management of COPD.

High FeNO level<sup>103</sup> and/or blood eosinophilia<sup>104,105</sup> have been identified as individual and/or surrogate markers of the response to steroids in patients with COPD-A. These findings support the view that ICS treatment may be beneficial in patients with COPD-A.

Table 6: Diagnosis of COPD-A:

| History and Symptoms  | Physical Findings                            | Investigations  |
|---|--|---|
| Phenotypes: Childhood Onset asthma / Adulthood onset asthma may have seasonal variation / Allergy to multiple Allergens | Prolonged Expiration<br>Polyphonic<br>Ronchi | Sputum Eosinophil $\geq 3/HPF$ <sup>106</sup>   |
| Recurrent variable Cough/SOB and/or wheeze  | Scattered Ronchi                             | Blood Total Eosinophil $\geq 300/\mu L$ <sup>107</sup>  |
| Variable amount of sputum   | Hyper inflated Lung ( $\pm$ )                | Blood IgE $\geq 100iu/\mu L$ <sup>108</sup>   |
| History of atopy e.g.; Allergic Rhinitis  | Crepitation infrequent                       | FeNO $\geq 20$ PPB <sup>100,101,109</sup>   |
| Family history of Bronchial Asthma  |  | Spirometry partially Reversible (not fit with asthma) <sup>110</sup> Experience of Ingenious *Pulmo-fit suggest that absolute ratio improvement of FEV <sub>1</sub> (in ml) / FVC (in ml) $> 1$ indicating asthma component more significant and absolute Improvement of FEV <sub>1</sub> (in ml) / FVC (in ml) $< 1$ indicating COPD component more important. |
| Smokers/exposure to smoke - Long time   |  | HRCT scan shows air trapping <sup>111</sup> but it did not show any strong association with the physiologic parameters of airway obstruction, either FEV <sub>1</sub> or FEF <sub>25-75</sub> .   |
|   |  | Skin prick test positive to multiple triggers including House Dust Mites and pollens <sup>112</sup>   |

**NOTE: Symptoms / History, physical findings and at-least 2 positive investigation findings along with post-Bronchodilator FEV<sub>1</sub>/FVC% <70% of predicted diagnostic of COPD-A.**

\*If sputum Eosinophil plenty or IgE ≥ 400 IU or blood Eosinophil ≥ 600/ml or FeNO ≥ 40 ppm, any single criteria of investigation may suggestive of COPD-A. Summation of more and more criteria confirm diagnosis.

\*Pulmo-fit suggest that absolute ratio improvement of FEV<sub>1</sub>(in ml) / FVC (in ml) > 1 indicating asthma component more significant and < 1 indicating COPD component more important.




Figure 17: Calculation of absolute ratio of improvement

## 9.2 COPD and Bronchiectasis: COPD-B

In moderate to severe COPD, prevalence of bronchiectasis varies from 57% to 64%<sup>9,113</sup>. Bronchiectasis independently increases risk of all cause of morbidity and mortality in patients with moderate to severe COPD<sup>114,115</sup>. This sub-class of COPD is more prone to infective exacerbations<sup>46</sup>. COPD patients with bronchiectasis (COPD-B) can be diagnosed by clinical history, sputum production more than 5ml (1 TSF) per-day, persistent inspiratory crepitations along with chest x-ray findings with or without HRCT Scan chest findings are important to sub classify the disease<sup>116</sup>. In chest x-ray persistent line shadow and/or inhomogeneous opacity not cleared by proper course of antibiotic, with or without ring shadows or cystic shadows in a patient with COPD may be a clue for diagnosis of COPD with bronchiectasis (COPD-B)<sup>117</sup>.

Table 7: Diagnosis of COPD-B

| History and Symptoms   | Physical Findings  | Investigations   |
|--|--|--|
| Symptoms after 40 years of age   | Usually near normal size Lung  | Sputum Neutrophilia <sup>118</sup>   |
| No history of atopy/ Allergy   | Prolonged Expiration   | Blood Total Eosinophil < 300/ $\mu$ L <sup>106</sup>                           |
| Long time Smokers / exposure to smoke  | Scattered crepitations more in basal   | Blood IgE <100iu/ $\mu$ L <sup>108</sup>                                       |
| Less variable Symptoms, usually progressive and more persistent Symptoms                                     | Scattered Ronchi   | FeNO low, <20 PPB <sup>100,101,109</sup>                                       |
| Frequent mucoid, mucopurulent or purulent sputum   |  | Spirometry: usually not Reversible   |
|  |  | CXR and or HRCT scan shows multiple cystic or tram line shadows <sup>119</sup> |
| Severity of Bronchiectasis Should be measured by FACED Score:<br>Mild: 1-2;<br>Moderate: 3-4;<br>Severe: 5-7 | Severity of score should be measured by Modified FACED SCORE <sup>120</sup> .<br>The FACED score consists of five dichotomized variables:<br>F – FEV <sub>1</sub> ( $\geq$ 50% = 0, < 50% = 2)<br>A – Age (< 65 years = 0, $\geq$ 65 years = 2)<br>C – Chronic colonization (no Pseudomonas = 0, presence of Pseudomonas =1 )<br>E – Extension (1 – 2 lobes affected = 0, > 2 lobes affected= 1)<br>D – Dyspnea – mMRC 0 – 2 = 0, mMRC 3 – 4 = 1 |  |

NOTE: For Bangladeshi Patient: we used 65 years cut off point for age instead of 70 years, which is in original FACED score. FACED score is important for diagnosis of severity of bronchiectasis in COPD.

Symptoms, physical findings, X-Ray Chest and or-HRCT scan chest and any 2 investigation findings of Table 7 along with post-Bronchodilator FEV<sub>1</sub>/FEV <0.7 is important for diagnosis of suggestive of COPD-B.

If sputum Neutrophil plenty or IgE less than 50 IU or blood Eosinophil less than 100/ $\mu$ L with FeNO less than 10 PPB: Presence of more and more criteria suggestive of COPD B.

HRCT scan of chest is important not only for diagnostic confirmation of COPD-B, but it is also an essential helping tool to rehabilitate the patient properly. Normally the lower respiratory tract is free from any microorganism. In stable COPD patients, sputum neutrophils and IL8- levels are higher than those in healthy subjects, which suggests ongoing neutrophilic inflammation in the airways<sup>121</sup>. This inflammation along with recurrent flare-up of infection, associated endobronchial obstruction by secretion and bronchoconstriction leads to development of bronchiectatic changes in COPD patients<sup>15</sup>.

Recurrent COPD exacerbations are associated with a heightened airway inflammatory burden, and the presence of lower airway bacterial colonization, which in turn has been shown to be an independent stimulus to airway inflammation in COPD<sup>122</sup>. In addition, it is now clear that lower airway bacterial colonization in the stable state is associated with exaggerated symptoms and sputum purulence at exacerbation<sup>123</sup>. The possible role of unrecognized bronchiectasis in orchestrating such relationships in COPD has not been previously assessed<sup>124</sup>.

HRCT is now accepted as the imaging modality of choice for the evaluation of bronchiectasis and emphysema<sup>125</sup>. Thin-section CT has been shown to have discriminatory value in obstructive lung disease<sup>128</sup>. However, there is no consensus to date on the role of HRCT in quantifying the structural changes of bronchiectasis in patients with COPD, and its use may have had a number of limitations. Previous studies of HRCT scanning in patients with clinical bronchiectasis<sup>126</sup> and one study of patients with cystic fibrosis, found significantly higher mean bronchiectasis scores<sup>127</sup>. The extent of bronchiectasis has been shown to be negatively correlated with FEV<sub>1</sub>% predicted<sup>128</sup> suggesting that in patients with COPD and bronchiectasis may develop more features of progressive airway obstruction. Severity of Bronchiectasis Should be measured by FACED Score (Table 7), which is better than Chest X-Ray, HRCT or FEV<sub>1</sub> alone to measure extent of bronchiectasis.

There are a number of possible reasons why bronchiectasis is detected most frequently in the lower lobes. In a previous study of patients with chronic purulent sputum production predominantly lower lobe distribution of bronchiectasis was found in subjects with impaired muco-ciliary clearance, one of the impaired host defense mechanisms seen in COPD<sup>129</sup>.

A lower lobe distribution is most often seen in patients with a history of childhood viral infections, a suggested risk factor for COPD<sup>4,9,15</sup>. It is possible that multiple physiological and pathologic alterations, including damaged mucociliary transport, localized or diffuse peripheral obliteration of the bronchial tree or lung tissue scarring are found in COPD<sup>130</sup>. In the context of an already disrupted lung parenchyma, produced structural changes of bronchiectasis are seen on HRCT<sup>124</sup>. It is important to establish criteria for the detection of these structural changes and their significance in COPD and for clarifying how they may relate to the natural history of this condition<sup>131</sup>.

A high prevalence of bronchiectasis has been demonstrated in an unselected group of patients with a primary care diagnosis of COPD<sup>132</sup> and studies of patients with Alpha<sup>1</sup> antitrypsin deficiency disease have suggested that bronchiectasis may be present either concomitantly before 50 years of age, and bronchiectasis may be developed before the development of emphysema<sup>133</sup>. The Alpha<sup>1</sup> antitrypsin status of all patients are not estimated. However, it is possible that some of these patients had bronchiectasis and then developed COPD with emphysema in addition to this at a later age<sup>113</sup>.

Management of COPD-B is mainly control of infection, prevention of infection flare up along with sustained bronchodilation<sup>139</sup>. Azithromycin alternate day or thrice a week throughout the year may prevent exacerbations<sup>134</sup>. Tobramycin nebulization<sup>135</sup>, influenza<sup>136</sup>, pneumonia vaccine<sup>137</sup>, sublingual lyophilized bacterial lysate may decrease recurrent infective exacerbations<sup>138</sup> in COPD-B patient.

### 9.3 COPD and GERD: COPD-G

The prevalence of gastro-esophageal reflux (GERD) was about 26.7% in COPD patients<sup>9</sup>. According to the report of Casanova et al<sup>139</sup> GERD was about 62% confirmed by esophageal 24-hour PH monitoring but 58% of them did not have any reflux symptoms. That means COPD patient with symptomatic GERD about 25-35%. Remaining 30-35% COPD have asymptomatic GERD with COPD. The annual rate of exacerbation of COPD is about two times higher in patient with COPD with GERD (COPD-G) compared to those without GERD symptoms<sup>9,15</sup>. Aspiration into airway in GERD patients can trigger exacerbation of COPD and enhance inflammation and induces pulmonary fibrosis<sup>9</sup>. Identification of GERD subgroup of COPD (COPD-G) is very important for prevention of progression of disease (Table 8).

An explanation for the association between GERD and exacerbations of COPD could be that the aspiration of gastric acid causes airway inflammation<sup>140</sup>. Proton pump inhibitors virtually abolish acid secretion in normal clinical doses<sup>141</sup>.

There is a null- association between GERD and COPD in users of acid inhibitory treatment, which altogether suggest that the acidity of the reflux content could be the key to the link between reflux and COPD exacerbations<sup>142</sup>. There are two ways for exacerbations of COPD. (A) Reflux theory: a direct mechanism by which micro-aspiration of gastro-duodenal contents can cause damage to the pulmonary tree. (B) Reflex theory: an indirect mechanism in which distal esophageal reflux stimulates the vagal nerve which on its turn leads to bronchoconstriction<sup>143</sup>. Furthermore, treatment with regular acid inhibitory treatment in COPD is symptomatic treatment and not without adverse effects. In fact, previous studies showed that the use of proton-pump inhibitors can be associated with an increased risk of pneumonia<sup>144,145</sup>. In addition, the effect of GERD on exacerbations could very well be influenced by other factors such as nonacidic reflux and pepsin<sup>146</sup>.

Explanation linking GERD with exacerbations is that GERD could cause symptoms such as cough that could be perceived<sup>147</sup> as an exacerbation by both patients and doctors, thereby resulting in the treatment with systemic corticosteroids. Treatment for GERD-related cough, using acid inhibitory treatment regularly, may reduce this symptom sufficiently to prevent treatment with systemic corticosteroids<sup>148,149</sup>. In fact, as in previous studies it was observed that individuals with COPD and GERD report significantly more breathlessness, wheezing and chronic bronchitis and have more often a history of respiratory infections than individuals without GERD<sup>140</sup>. Proton pump inhibitor therapy, anti-reflux therapy, change of food habit and sustained bronchodilation are very important to control COPD-G patient<sup>150</sup>.

HRCT scan may show basal fibrosis may mimic Idiopathic Pulmonary Fibrosis pneumonia (IPF) with or without esophageal dilatation<sup>151,152</sup>. Basal lung fibrosis with or without esophageal dilatation may a very good marker of GERD in COPD patient.

Table 8: Diagnosis of COPD-G

| 1 | Symptoms   | Physical Findings                  | Investigations                          |
|---|--|------------------------------------|---|
| 2 | Symptoms after 40 years  | Usually near normal size Lung      | Sputum Neutrophilia                     |
| 3 | Heartburn, Regurgitation, Dysphagia (HRD) score (Table 9) suggestive of GERD | Recurrent wheeze after taking food | Blood Total Eosinophil less than 300/ML |



|   |  |  |  |
|---|--|--|--|
| 4 | Long time Smokers/exposure to smoke  | Basal Crepitations like Lung fibrosis or ILD | Blood IgE less than 100/ML                           |
| 5 | Coughing and shortness of breath after taking food or drinks   | Ronchi after taking food                     | FeNO very low, at least less than 20 PPB             |
| 6 | Coughing and Shortness of breath few hours after sleep   |  | Esophageal pH monitoring is important <sup>140</sup> |
| 7 | Basal fibrosis in HRCT scan may mimic Idiopathic Pulmonary Fibrosis (IPF) with or without esophageal dilatation. Esophageal dilatation (ED) <sup>153</sup> is defined as the presence of air bubbles greater than 10 mm in the supra ventricle (SV) and ventricle (CV), and air bubbles >15 mm at the ventricle to the lower esophageal sphincter (V-LES). |  |  |

Table 9: HRD score Uploaded by Rachel Mary Thomson

| Symptom                  | Description   | Score |
|--------------------------|---|-------|
| <b>Heartburn (H)</b>     | None  | 0     |
|                          | Occasional, brief episodes; readily controlled by simple antacids (e.g, Liquid Antacid  | 1     |
|                          | Frequent (>2/week) episodes causing moderate discomfort & requiring medication prescribed by your doctor for relief                                   | 2     |
|                          | Daily attacks causing interference with work/activities; night time attacks interfering with sleep; only controlled by prescription medication        | 3     |
| <b>Regurgitation (R)</b> | None  | 0     |
|                          | Occasional episodes; mostly after meals & not predictable   | 1     |
|                          | More frequent attacks; predictable by posture   | 2     |
|                          | Daily episodes interfering with life: severe night time episodes causing choking coughing or the sensation of acid/food/fluid is going into the lungs | 3     |

|                  |  |     |
|------------------|--|-----|
| Dysphagia<br>(D) | None   | 0   |
|                  | Occasional transient sensation of food “sticking” in the gullet  | 1   |
|                  | Episodes of food getting stuck requiring liquids to clear  | 2   |
|                  | Progressive difficulty swallowing solids requiring medical attention/admission to hospital. Need for gastroscopy/telescopic examination of the stomach/gullet. | 3   |
|                  | Total  | 0-9 |

The DeMeester symptom score for GERD<sup>154</sup>. 4 or less was considered mild GERD, scores between 5-6 were regarded as moderate GERD, and a score 7-9 was regarded as severe GERD. Due to pulmonary fibrosis patient may have  $FEV_1/FVC\% >0.7$  due to concomitant restriction.




Figure 18: CT Scan at upper and lower level of chest

NOTE: Dilated esophagus showing in CT SCAN of chest at the level of upper chest and lower chest<sup>153</sup>.

Symptoms, physical findings HRD score 4 or more and Esophageal pH monitoring (if available findings) -along with post-Bronchodilator  $FEV_1/FVC$  Ratio 0.7 or less is important for diagnosis of suggestive of COPD-G.

#### 9.4 COPD and Obstructive Sleep Apnea (OSA): COPD-0

When obstructive sleep apnea (OSA) and chronic obstructive pulmonary disease (COPD) coexist in the so-called “overlap” syndrome, a high risk for mortality and morbidity has been reported. There is high prevalence (29% to 65.9%) of Obstructive Sleep Apnea in Patients with Moderate to Severe Chronic Obstructive Pulmonary Disease<sup>155,156</sup>. COPD alone can cause subjective and ob-

jective changes during sleep. When those with either chronic bronchitis or emphysema were surveyed across a broad range of symptoms, “sleep difficulties” were endorsed as occurring “almost always” or “always” in 43% of subjects (third most common, after dyspnea and fatigue)<sup>157,158</sup>. Specifically, patients with COPD report more difficulty in both initiating and maintaining sleep than controls, and also complain of excessive daytime sleepiness<sup>159</sup>. Sleep architecture in some of these same patients was notable for many arousals. More than just the diagnosis of COPD, the presence of COPD symptoms such as cough or sputum production or wheezing strongly correlated with difficulty falling or staying asleep<sup>160,161</sup>. Other investigations have objectively confirmed poor sleep quality, with decreased total sleep time and decreased sleep efficiency<sup>96,162</sup>. (Table 10).

#### **9.4.1 Sleep and Breathing<sup>162</sup>**

A brief review of the normal changes in respiration that occur with sleep onset and the various sleep stages is helpful to understand the changes that occur during sleep in those with COPD. In normal subjects, minute ventilation drops from wakefulness to non-rapid-eye-movement (non-REM) sleep, and drops further during REM sleep (about 15%, compared to the awake value). Most of the drop in minute ventilation is due to a decrease in tidal volume that is not fully compensated by a concomitant increase in respiratory rate. There is a blunted ventilatory response to hypoxia and hypercapnia, again with the greatest changes during REM sleep. REM is characterized by skeletal muscle atonia, except for the diaphragm, and shallow, irregular breathing. Finally, even in normal subjects without OSA, upper-airway resistance increases during sleep<sup>163</sup>.

#### **9.4.2 Nocturnal Oxygen Desaturation<sup>155</sup>**

The most significant sleep abnormality associated with COPD is nocturnal oxygen desaturation. Even without any upper-airway contribution, various studies have reported that 27–70% of patients with COPD with awake oxygen saturation of 90–95% can experience substantial desaturation at night, particularly during REM sleep<sup>164</sup>. Nocturnal oxygen desaturation can be defined or measured in terms of oxygen nadir or time below some oxygen saturation limit, such as 88% or 90%. The desaturation nadir is more profound than during exercise, with oxygen saturation falling an average of  $6 \pm 4\%$  during peak exercise and  $13 \pm 9\%$  during sleep<sup>165</sup>. Awake oxygen saturation has the greatest predictive value, although it imperfectly predicts nocturnal desaturation<sup>166</sup>. Daytime PCO<sub>2</sub> has also been found to be predictive. Perhaps most clinically relevant, nocturnal oxygen desaturation is a marker of increased mortality in COPD<sup>167</sup>.

American Thoracic Society/European Respiratory Society guidelines also suggest that those with relatively mild COPD and evidence of pulmonary hypertension should be referred for overnight testing<sup>168</sup>. Daytime hypoxemia with or

without hypercapnia and pulmonary hypertension in patients known to have only one disease (either OSA or COPD), whatever in severity, should prompt assessment for the other disorder<sup>162</sup>. CPAP therapy is an effective therapeutic option in the majority of patients with obstructive sleep apnea, even if severe, and with normal awake respiratory function<sup>169</sup>. For the subset of patients with OSA associated to COPD, especially when hypercapnia is present in whom CPAP may be ineffective and/or not tolerated.; BiPAP may be an effective and well-tolerated treatment modality for them<sup>170,171</sup>.

Table 10: Diagnosis of COPD-O

| Symptoms  | P/E                           | Investigations  |
|---|-------------------------------|---|
| Symptom after 40 years  | Usually near normal size Lung | ABG showing Hypoxemia i.e. PO <sub>2</sub> less than 80mm of Hg or 10.6 mmol <sup>172,173</sup> |
| Excess snoring during sleep and day time sleepiness   |                               | PASP >30mm <sup>174</sup>   |
| Long time Smokers/ exposure to smoke  |                               | Fall of SpO <sub>2</sub> >4% during sleep <sup>175</sup> .                                      |
| Day time Feelings of Shortness of breath, malaise and easy fatigability   |                               | Sleep test is most important tool to diagnose COPD-O <sup>155</sup>                             |
| <b>Chest radiography Pulmonary Arterial:</b> Hypertension (PAH) can be diagnosed with 98% sensitivity, if the diameter of the right descending pulmonary artery (PA) is increased (>16 mm, posteroanterior projection) together with an increased diameter of the left descending PA (>18 mm, left lateral projection) <sup>176</sup> |                               |   |
| HRCT scan CHEST may show prominent vascular markings. The typical radiographic appearance described is that of a diffuse bilateral reticulonodular pattern associated with enlarged central pulmonary arteries <sup>50,177</sup>  |                               |   |
| Echocardiography: As non-invasive tool for the diagnosis of PAH in COPD. Color Doppler transthoracic echocardiography is the most important diagnostic test used for PAH in COPD <sup>178-180</sup> .   |                               |   |

NOTE: Symptoms, physical findings and any 2 investigation findings suggestive of COPD-O.

Sleep test is the gold standards test for OSA along with post-Bronchodilator FEV<sub>1</sub>/FVC <0.7 or less is important for diagnosis of COPD-O. Due to Obesity patient may have FEV<sub>1</sub>/FVC % >0.7 due to concomitant restriction.

## 9.4.3 Screen for Obstructive Sleep Apnea (OSA):

### 9.4.3.1 Epworth Sleepiness Scale

The Epworth Sleepiness Scale is used to diagnose obstructive sleep apnea (OSA)<sup>181</sup>.

If you suspect you might have OSA, answer the following questions and take your answers to GP.

Description of Epworth Sleepiness Scale:

How likely are you to fall asleep in the following situations, in comparison to feeling just tired?

This refers to your usual way of life in recent times.

Even if you haven't done some of these things recently, try to work out how they would have affected you.

Use the following scale to choose the most appropriate number for each situation:

- 0 = would never doze
- 1 = slight chance of dozing
- 2 = moderate chance of dozing
- 3 = high chance of dozing

It's important that you answer each question as best you can.

Table 11: Epworth Sleepiness Scale

| Situation   | Chance of dozing<br>(0 = low, 3 = high) |
|---|---|
| Sitting and reading   | 0 to 3                                  |
| Watching TV   | 0 to 3                                  |
| Sitting still in a public place (e.g. a theatre, a cinema or a meeting) | 0 to 3                                  |
| As a passenger in a car for an hour without a break                     | 0 to 3                                  |
| Lying down to rest in the afternoon when the circumstances allow        | 0 to 3                                  |
| Sitting and talking to someone  | 0 to 3                                  |
| Sitting quietly after lunch without having drunk alcohol                | 0 to 3                                  |
| In a car or bus while stopped for a few minutes in traffic              | 0 to 3                                  |
| TOTAL   | 0 to 24                                 |

### 9.4.3.2 What does the score mean?

Scores can be interpreted as follows:

- 0-5 lower normal daytime sleepiness
- 6-10 normal daytime sleepiness
- 11-12 mild excessive daytime symptoms
- 13-15 moderate excessive daytime symptoms
- 16-24 severe excessive daytime symptoms

Table 12: STOP-BANG Questionnaire

It is another Practical Approach to Screen for Obstructive Sleep Apnea<sup>182,183</sup>. STOP-Ban questionnaire is a useful tool for screening OSA (Fig. 19)

| <b>STOP-Bang questionnaire</b>  |   |
|---|---|
| Please answer the following questions by checking “yes” or “no” for each one.                     |   |
|   | <b>Yes    No</b>                                  |
| Snoring (Do you snore loudly?)  | <input type="checkbox"/> <input type="checkbox"/> |
| Tiredness (Do you often feel tired, fatigued, or sleepy during the daytime?)                      | <input type="checkbox"/> <input type="checkbox"/> |
| Observed Apnea (Has anyone observed that you stop breathing, or choke or gasp during your sleep?) | <input type="checkbox"/> <input type="checkbox"/> |
| High Blood Pressure (Do you have or are you being treated for high blood pressure?)               | <input type="checkbox"/> <input type="checkbox"/> |
| BMI (Is your body mass index more than 35 kg per m <sup>2</sup> ?)                                | <input type="checkbox"/> <input type="checkbox"/> |
| Age (Are you older than 50 years?)  | <input type="checkbox"/> <input type="checkbox"/> |
| Neck Circumference (Is your neck circumference greater than 40 cm [15.75 inches]?)                | <input type="checkbox"/> <input type="checkbox"/> |
| Gender (Are you male?)  | <input type="checkbox"/> <input type="checkbox"/> |

*Score 1 point for each positive response.*  
*Scoring interpretation: 0 to 2 = low risk, 3 or 4 = intermediate risk, ≥5 = high risk.*

Source: University Health Network, Toronto, Ontario, Canada ([www.stopbang.ca/osa/screening/php](http://www.stopbang.ca/osa/screening/php)). Used with permission from Sauk Prairie Healthcare.

Figure 19: STOP-Bang questionnaire

## 9.5 Mixed Disorder:

When multiple coexisting illnesses develop and perpetuated during the progress and stage change of COPD, amalgamated symptoms, signs and investigation findings, are sometimes difficult to differentiate various coexisting illnesses. In those cases, clinical judgement of each component is important for pin point diagnosis and for management plan of COPD.

## 10 CHAPTER 10: COMMON PROBLEM FOR COPD

### 10.1 Major triggers, common to all COPD patients:

#### 10.1.1 Air Pollution<sup>9</sup>:

Air pollutants such as particulate materials (PM) from fossil, diesel or octane combustion can cause inflammation in the lung and further impairs the reduced pulmonary function in COPD patients<sup>184-186</sup>. When exposed to particle pollution, patients with COPD usually have more emergency room visit, hospital admission and/or even death in some cases. Infection is one of the inducing factors of exacerbations of COPD. As particulate materials can bring many micro-organisms on the surface, inhalation of such materials may contribute to more frequent infective exacerbation of COPD also. Other mechanisms including the impairment of mucociliary clearance, increased adherence of virus to respiratory mucus cells, and impairment of the resistance ability of immune system are all involved in the adverse effects of pollutants<sup>187</sup>.

Several studies link outdoor air pollution to increased risk of COPD exacerbations. The Air Pollution and Health, a European Approach (APHEA) project analyzed data from 6 European cities and found increased risk of COPD hospital admissions with several air pollutants, including NO<sub>2</sub>, O<sub>3</sub>, sulfur dioxide (SO<sub>2</sub>) and black smoke<sup>188</sup>. A study of PM<sub>10</sub> and hospital admissions for COPD found a 2.5% increase in admissions for every 10µg/m<sup>3</sup> increase in PM. In addition, a meta-analysis of 18 studies of PM<sub>10</sub> and exacerbations found a 10µg/m<sup>3</sup> increase in daily PM<sub>10</sub> was associated with a 2.7% increase in COPD hospitalizations<sup>189</sup>. The data linking air pollution with COPD exacerbations for O<sub>3</sub> is also convincing as higher O<sub>3</sub> concentrations have been shown to increase hospital and emergency department visits for lower respiratory disease, including COPD<sup>190</sup>.

#### 10.1.2 Temperature

Extremes of temperature—both heat and cold—have been associated with increased respiratory morbidity in COPD<sup>190</sup>.

Studies have consistently found that elderly individuals and those with underlying cardiac and respiratory diseases, including COPD, are at increased risk for adverse health effects of heat exposure<sup>191</sup>. For example, a study across 12 European cities estimated that the effect of hot temperatures during summertime can increase the risk of death attributable to COPD by as much as 25%<sup>192</sup>. A study in New York City found that the risk of COPD hospitalization increased by 7.6% for every 1°C increase above a threshold temperature of 29°C<sup>9,193</sup>. A large study in Taiwan using national health insurance registry data detected a 0.8% increase in COPD exacerbations for every 1°C decrease in mean daily temperature<sup>194</sup>. Bronchoconstriction and inflammation that may occur in the setting of cold exposure<sup>195</sup>. Recent evidence suggests a role for mucous hypersecretion as a potential mediator of the COPD response to cold temperature<sup>196</sup>.

### 11.1 Comorbidities<sup>52</sup>:

Comorbidities such as cardiac disease, diabetes mellitus, hypertension, osteoporosis, and psychological disorders are commonly reported in patients with chronic obstructive pulmonary disease (COPD) but with great variability in reported prevalence<sup>197,198</sup>. The most common comorbidities differ between men and women. Men have more IHD and stroke than women. Women are more likely to demonstrate anxiety, depression and Osteoporosis than men<sup>199</sup>.

### 11.2 Cardiac Disease

COPD patients possess an increased burden of Ischemic heart disease, cardiac arrhythmias and heart failure when compared to the normal population<sup>200,201</sup>. Ischemic Heart Disease (IHD) is an important cause of hospitalization and mortality in COPD, even affecting those with mild COPD. In addition to the high separate prevalence of COPD and IHD, these conditions share same risk factors, e.g. advanced age, smoking, low socioeconomic condition and sedentary lifestyle. Systemic inflammation, oxidative stress, hypoxia particularly hypoxic spell during sleep, acidosis, bacterial colonization and haemo-dynamic derangements are likely to be added contributing factors<sup>122</sup>. Independent of smoking and other risk factors, impaired lung function with or without hypoxemia is a major risk factor for IHD and arrhythmia (with or without hyperlipidemia), with the relationship being strongest for fatal cardiac events<sup>202</sup>. In COPD, arterial stiffness increases during exacerbation and is associated with COPD severity (measured as airflow limitation or degree of emphysema), inflammation, oxidative stress and sympathetic nervous system (SNS) tone, which increases risk of acute ischemic cardiac events<sup>130</sup>. It is now well documented that a statistically significant improvement in ischemic heart disease after standard pulmonary rehabilitation and after treatment with combination ICS/LABA or LAMA as well as possible improvement with supplemental oxygen<sup>203</sup>.

Ingenious Pulmo-Fit, Dhaka Registry demonstrated that in total 2551 cases tele-cardiac ECG monitoring done in 2019. Out of 2551, 1652 (64.8%) patients were COPD stage 1 to 4 and 899 (35.2%) patient were Non-COPD. COPD patients with tele-cardiac monitoring during Modified Cardio-Pulmonary Exercise (M-CPET) demonstrated more arrhythmias than those without COPD; atrial fibrillation/flutter or atrial ectopic were identified in 426(16.7%) versus 107 (4.2 %) (OR 2.572 95% CI 2.043 - 3.237), and non-sustained Ventricular extra systole to non-sustain ventricular tachycardia in 136 (8.2%) versus 24(2.7%) (OR 3.271 95% CI 2.103 - 5.088).



This study population of Ingenious Pulmo-Fit, Dhaka might be highly selected group, which potentially limits the broad application of the results. However, one may understand potential arrhythmias may develop during exercise of COPD patients. These Cardiac arrhythmias sometimes may be an important obstacle during pulmonary rehab program. If palpitations continue and are so bothersome that they prevent patients from exercising, then the appropriate next step is to look into fixing the problem. Many arrhythmias can be treated or cured with medications<sup>204</sup>.

Medications used in the treatment of COPD also have potential to impact on cardiac morbidity and mortality, due to intrinsic effects on chronotropy and muscle action potentials or due to side effects such as hypokalemia<sup>205</sup>. Medications concerned include beta-agonist and antimuscarinic bronchodilators and methylxanthines. More recently, macrolide antibiotics, which in chronic dosing have been shown to reduce respiratory exacerbations, have been added to the list, due to an association with QT prolongation and bradycardia<sup>206</sup>. Randomized controlled trials (RCT) of chronically dosed azithromycin have not demonstrated adverse cardiac effects in the clinical setting<sup>207</sup>, particularly when known drug interactions are avoided. Likewise, for most inhaled bronchodilators, when used at therapeutic dose in stable COPD, there are no proven adverse effects on safety<sup>3</sup>.

### 11.3 Heart Failure

Chronic obstructive pulmonary disease (COPD) and heart failure (HF) are both common diseases with major impact and seem to coexist more frequently than expected from their distinct population prevalence<sup>208,209</sup>.

The natriuretic peptides, including BNP and NT-pro BNP, can assist in identifying heart failure in the setting of acute breathlessness, but do not exclude comorbid COPD, but currently it has an unclear diagnostic role in stable disease<sup>210</sup>. The prevalence of heart failure in COPD patients is estimated at 20 to 32%. For the converse situation in heart failure, COPD prevalence has been previously quoted as 10 to 33%<sup>63</sup>.

#### 11.3.1 Role of BNP and NT-proBNP in clinical practice<sup>211</sup>

B-type natriuretic peptide, which is also called brain-type natriuretic peptide (BNP), was first described in 1988 after isolation from swine brain. However, it was soon found to originate mainly from the heart, representing a cardiac hormone. BNP belongs to the natriuretic peptide family together with other structurally similar peptides, namely atrial natriuretic peptide (ANP), C-type natriuretic peptide (CNP), and urodilatin. The natriuretic peptides have in common a characteristic biochemical structure which consists of a 17 amino acid ring and a disulfide bridge between two cysteine molecules. The major source

of BNP synthesis and secretion is the ventricular myocardium. Whereas ANP is stored in granules and can be released immediately after stimulation, only small amounts of BNP are stored in granules and rapid gene expression with de novo synthesis of the peptide is the underlying mechanism for the regulation of BNP secretion. BNP is synthesized as a prohormone (pro-BNP) comprising 108 amino acids. Upon release into the circulation it is cleaved in equal proportions into the biologically active 32 amino acid BNP, which represents the C-terminal fragment, and the biologically inactive 76 amino acid N-terminal fragment (NT-proBNP). Both molecules are constantly released and can be detected in the blood. The main stimulus for increased BNP and NTproBNP synthesis and secretion is myocardial wall stress. Furthermore, factors such as myocardial ischemia and endocrine (paracrine) modulation by other neurohormones and cytokines are also of importance. In the systemic circulation BNP mediates a variety of biological effects by interaction with the natriuretic peptide receptor type A (NPR-A) causing intracellular cGMP production. The physiological effects of BNP are manifold and comprise natriuresis/diuresis, peripheral vasodilatation, and inhibition of the renin–angiotensin–aldosterone system (RAAS) and the sympathetic nervous system (SNS). BNP is cleared from plasma by binding to the natriuretic peptide receptor type C (NPR-C) and through proteolysis by neutral endopeptidases. In contrast, NT-proBNP is mainly cleared by renal excretion. However, recent studies suggest that there might also be other important clearing mechanisms for NTproBNP. The half-life of BNP is 20 mins whereas NT-proBNP has a half-life of 120 mins, which explains why NT-proBNP serum values are approximately six times higher than BNP values, even though both molecules are released in equimolar proportions. Cut-off values of BNP<sup>212</sup> and NTproBNP<sup>213</sup> for the diagnosis of heart failure of patients presenting with dyspnea.

Table 13: Cut-off values of BNP and NTproBN

|   | HF likely | HF likely |
|---|-----------|-----------|
| BNP (pg/ml)                               | 100       | 500       |
| NT-proBNP (pg/ml), age <50 years          | 300       | 450       |
| NT-proBNP (pg/ml), age more than 50 years | 300       | 900       |

The diagnostic performance of BNP and NT-proBNP is comparable and there is no meaningful difference between them. They reflect hemodynamic myocardial stress independent of the underlying pathology, thus they are not specific for a distinct pathology of heart failure. Their particular strength is to rule out heart failure in patients presenting to the emergency department with shortness of breath.

Table 14: Diagnosis of COPD and Heart Failure

| History and Symptoms   | Physical Findings  | Investigations   |
|--|--|--|
| Age usually 60 or more and post bronchodilator FEV <sub>1</sub> /FVC and/or FEV <sub>1</sub> evidence of COPD                                    | Prolonged Expiration   | BNP <sup>212</sup> or NT-proBNP <sup>213</sup> = Rule in HF (Table 13)   |
| History of old MI/ Angina like chest pain during exertion/ PND/ Orthopnea  | Ronchi and bilateral crepts  |  |
| Persistent and/or disproportionate shortness of breath and marked limitation of activities despite taking adequate Dual bronchodilator LABA/LAMA | Hyper inflated Lung and Apex beat may be shifted due to cardiomegaly | Color doppler <sup>214</sup> <ul style="list-style-type: none"> <li>• Hyperdynamic= LVEF greater than 70%</li> <li>• Normal = LVEF 50% to 70% (midpoint 60%)</li> <li>• Mild dysfunction = LVEF 40% to 49% (midpoint 45%)</li> <li>• Moderate dysfunction = LVEF 30% to 39% (midpoint 35%)</li> <li>• Severe dysfunction = LVEF less than 30%</li> </ul> |
| History of Smoking and or smoke exposure<br>-Long time   |  | Color doppler ECHO <sup>50</sup><br>PAH PASP ≥ 35 mm of Hg Evidence of TR  |
| Presence of risk factor of IHD e.g. Hyperlipidemia, Diabetes and/or Hypertension <sup>215</sup>  |  |  |

NOTE: Symptoms / History, physical findings and any one positive investigation findings suggestive of COPD-and Heart Failure. Post-Bronchodilator FEV<sub>1</sub>/FVC% <70% may not be diagnostic of COPD due to associated HF associated pulmonary edema may cause restriction (FEV<sub>1</sub>/FVC% > 70%).

Since COPD and heart failure present with similar symptoms and frequently do coexist, the clinical implication is that the opportunity for intervention will be missed unless both diagnoses are specifically sought using careful clinical assessment in conjunction with appropriately directed investigations<sup>63</sup>.

### 11.3.2 Use of Cardio-selective Beta Blocker in COPD<sup>63</sup>:

A number of observational studies also lead confidence to cardio selective betablocker prescribing in COPD patients<sup>216</sup>. There is no evidence to suggest that baseline Beta-blocker therapy reduces the respiratory benefits or increases the cardiovascular risk of inhaled long-acting Beta-agonists in patients with chronic obstructive pulmonary disease and heightened cardiovascular risk<sup>217</sup>.

### 11.4 Osteoporosis

Patients with COPD are at increased risk for fracture due to the disease itself, the use of high dose corticosteroids and coexisting risk factors such as hypogonadism (induced by corticosteroid therapy itself in high doses in men and women), immobilization reduced muscle mass and other factors<sup>218</sup>. These patients may have reduced bone mineral density (BMD) due to a reduction in bone formation and perhaps increased bone resorption, the latter being primarily due to the underlying disease itself. A systematic review described an overall mean prevalence of osteoporosis of 35.1% (range 9-69%) with increasing odds ratios for osteoporosis associated with lower FEV<sub>1</sub>, lower BMI and lower fat-free mass index<sup>219</sup>. Patients with vertebral compression fractures, visualized on a lateral chest x-ray, had more frequent admissions, longer length of hospital stay, and increased mortality in the two years after admission<sup>220</sup>. Evidence for fracture risk reduction is available for alendronate, risedronate, etidronate and parathyroid hormone<sup>221</sup>. Although calcium supplementation has not been demonstrated to reduce the risk of fracture in osteoporosis, a reduction in remodeling rate with some possible benefit in slowing bone loss is possible so calcium supplements are appropriate. In Bangladesh Vitamin D insufficiency is epidemic 85.5%<sup>222</sup>. Therefore, vitamin D level should be corrected with supplements in all cases of COPD.

### 11.5 Frailty and COPD<sup>63</sup>

Frailty is common in seniors and is characterized by diminished physiological reserves and increased vulnerability to stressors<sup>223</sup>. In other words, Frailty is a state of vulnerability to poor resolution of homeostasis after a stressor event and is a consequence of cumulative decline in many physiological systems during a lifetime. Frailty is a state of older patient where physically or mentally patient is vulnerable and less able to recover quickly after illness or a stressful event.

Criteria for Diagnosis of the Frailty Syndrome: Three or More of Following Symptoms<sup>224</sup>

Table 15: Criteria for Diagnosis of the Frailty Syndrome

|    |  |
|----|--|
| 1. | Weakness   |
| 2. | Slow walking speed   |
| 3. | Self-reported exhaustion                                   |
| 4. | Low level of physical activity i.e. Low energy expenditure |
| 5. | Unintentional weight loss                                  |

Table 16: FRAIL SYNDROME: QUESTIONNAIRE BASED DIAGNOSIS<sup>225</sup>

## THE FRAIL NON-DISABLED (FIND) QUESTIONNAIRE

| Domain                      | Question   | Answer   | Score    |
|-----------------------------|--|--|----------|
| <b>DISABILITY</b>           | <b>A</b> Have you any difficulties at walking 400 meters?  | a. No or some difficulties                                 | <b>0</b> |
|                             |  | b. A lot of difficulties or unable                         | <b>1</b> |
|                             | <b>B</b> Have you any difficulties at climbing up a flight or stairs?  | a. No or some difficulties                                 | <b>0</b> |
|                             |  | b. A lot of difficulties or unable                         | <b>1</b> |
| <b>FRAILTY</b>              | <b>C</b> During the last year, have you involuntarily lost more than 4.5 kg?   | a. No  | <b>0</b> |
|                             |  | b. Yes   | <b>1</b> |
|                             | <b>D</b> How often in the last week did you feel that everything you did was an effort or that you could not get going | a. Rarely or sometimes (2 times or less/week)              | <b>0</b> |
|                             |  | b. Often or almost always (3 or more times per week)       | <b>1</b> |
|                             | <b>E</b> Which is your level of physical activity?   | a. Regular physical activity (at least 2-4 hours per week) | <b>0</b> |
| b. None or mainly sedentary |  | <b>1</b>   |          |

*if  $A+B \geq 1$ , the individual is considered as "disabled"*

*if  $A+B=0$  and  $C+D+E \geq 1$  the individual is considered as "frail"*

*if  $A+B+D+E=0$  the individual is considered as "robust"*

Figure 20: Self-reported questionnaire tool for frailty<sup>226</sup>

Frailty affects older people and particularly those with chronic conditions such as COPD. The mechanism underlying increased frailty in COPD is likely to be multifactorial. COPD affects older adults in whom other health conditions are more prevalent. In addition, COPD is associated with inflammation that affects multiple body systems, increased exacerbations, as well as lifestyle factors such as smoking and reduced physical activity, all of which may increase risk of frailty<sup>227</sup>.

### 11.5.1 Management of Frailty:

Frailty affects one in four patients with COPD referred for pulmonary rehabilitation and is an independent predictor of program non-completion. In summary, frailty is common in COPD and associated with poorer health outcomes, hospital admissions and failure to complete pulmonary rehabilitation<sup>228</sup>. Measuring frailty is useful in COPD and may identify vulnerable patients and allow earlier interventions such as pulmonary rehabilitation to minimize the development and impact of frailty on patients and careers as well as health and social care services<sup>229</sup>.

Table 17: Treatments of Frailty Syndrome

| Treatment   | Potential effect  | Evidence   |
|---|---|--|
| Vitamin D   | Improve Balance and Strength                            | Systematic review and meta-analysis <sup>230</sup> .   |
| Vitamin and Carotenoid  | Improved Walk-Speed, Strength                           | Cross-sectional study <sup>231</sup>   |
| Fruit and vegetables <sup>232</sup> , Vit E <sup>233</sup> and Omega-3 fatty Acids <sup>234</sup> , Nutritional Supplementation Combined with Low-Intensity Rehab Exercise <sup>235</sup> | Improve systemic inflammation and improve lung function | A Randomised Case control study (CCS) <sup>232</sup><br>Animal model CCS <sup>233</sup><br>Prospective CCS <sup>234</sup><br>A prospective RT <sup>235</sup> |
| Creatine  | Increase Strength                                       | RCT <sup>236</sup>   |

|  |  |   |
|--|--|---|
| Testosterone+ Exercise <sup>237</sup><br>Nandrolone+ Exercise <sup>238</sup><br>(No effect without exercise program)<br>A dietary pattern with more phyto-nutrient-rich plant foods, tea, omega-3-rich deep sea fish, and other protein-rich foods such as shellfish and milk had a reduced prevalence of frailty <sup>239</sup> | Increase Strength                                | RCT <sup>237</sup><br>RCT <sup>238</sup><br>Cross-sectional <sup>239</sup>  |
| Beta-hydroxy-beta-methyl butyrate  | Increase Strength                                | A Systematic Review <sup>240</sup>  |
| Rehabilitation for Older People in Longterm Care <sup>241</sup><br>Pulmonary Rehabilitation and COPD <sup>242</sup><br>Physical frailty and pulmonary rehabilitation in COPD <sup>228</sup>  | Improve Strength, Balance and Increase endurance | Cochrane Review <sup>241</sup><br>A Systematic Review <sup>242</sup><br>A prospective cohort study <sup>228</sup> |

## 11.6 Falls in COPD<sup>63</sup>

Chronic Obstructive Pulmonary Disease (COPD) and other chronic respiratory issues can cause shortness of breath, which may cause feelings of dizziness, weakness, and fatigability<sup>243,244</sup>. Chronic cough in COPD may lead to cough syncope and that leads to fall<sup>245</sup>.

Patients with COPD were 55% more likely to have an incident record of fall than non-COPD subjects (adjusted HR, 1.55; 95% CI 1.50 to 1.59). The greater falls risk in patients with COPD needs special consideration during management of COPD<sup>246</sup>.

In fact, COPD patients in moderate-severe stages and especially those in exacerbation have a high risk of falls<sup>247</sup>.

For this reason, it's especially important for seniors with breathing issues to be incorporated in pulmonary Rehab program for prevention of accidental falls<sup>248</sup>.

## 11.7 Aspiration and COPD

Aspiration of food and liquid is very common with COPD.

Subjective swallowing symptoms seems to be a common problem in patients with stable COPD. This problem is seen in all stages of the disease, but is more common in symptomatic patients and in patients with lower physical capaci-

ty<sup>249</sup>. Up to 70% COPD patients difficulty in swallowing or dysphagia leads to aspiration<sup>250</sup>.

Patients with COPD exhibit disrupted coordination of the respiratory cycle with deglutition. Disrupted breathing–swallowing coordination could increase the risk of aspiration in patients with advanced COPD and may contribute to exacerbations<sup>251</sup>.

The risk factors of aspiration in COPD reported include dyspnoea, dysphagia, emphysema, weakness or incoordination of throat muscles, decreased throat sensitivity, and an impaired cough reflex<sup>252</sup>. Impairment of the swallowing function, also referred to as oropharyngeal dysphagia (OD), can lead to pneumonia, aspiration, COPD exacerbations and malnutrition<sup>253</sup>.

Silent aspiration has also been reported in those with COPD, which can complicate dysphagia detection and management<sup>254</sup>.

Dysphagia in COPD is thought to be due to the disrupted coordination of the exhale-swallow-exhale respiratory cycle during swallowing<sup>251</sup>. Dysphagia and aspiration risk can be determined from adequate history of patients and from their spouse or care givers and clinical swallow examination<sup>255,256</sup>.

Management for swallowing problem will be provided on an individual basis and may involve the following:

### 11.7.1 Eating strategies which may prevent dyspnea

- Sit up straight
- Clear the airways of mucus before eating
- If supplemental oxygen is used, make sure this is worn while eating
- Avoid eating large meals, instead eat small nutritious meals and snacks more frequently
- Avoid drinking with meals, can take sips of water to facilitate swallowing.
- Take time, Eat slowly and take small bites and sips
- Reduce talking and distraction during meals
- Make sure mouth is empty before talking the next mouthful.
- Choose softer foods that are easier to chew and swallow, e.g. mashed boiled rice, potato, soups, bananas
- Limit foods that can cause bloating, e.g. beans, onions, cauliflower, soft drinks
- Rest for at least 20-30 minutes after eating in an upright position
- In patients who are underweight, protein and calorie intake can be boosted using high energy, nutrient-rich foods that are easily accessible, such as milk powder, cheese, cream, custard, peanut butter and milkshakes or a nutritionally complete oral supplement



Management for dysphagia and aspiration will be provided on an individual basis and may involve the following:<sup>257</sup>

- Rehabilitation exercises
- Swallowing – breathing retraining (compensatory swallowing techniques)
- Texture modification of diet and fluids
- Postural strategies
- Safe swallowing strategies

### 11.8 Lung cancer and chronic obstructive pulmonary disease:

Lung cancer is a serious health problem for Bangladesh. The first leading causes are in males are lung cancer and in females are breast cancer<sup>258</sup>. The risk of lung cancer in people who have pre-existing lung disease has been studied using case- control studies, which found an increased risk of lung cancer in people with bronchitis and emphysema, even after correcting for the smoking history<sup>259</sup>. Chronic obstructive pulmonary disease (COPD) and lung cancer are devastating pulmonary diseases that commonly coexist and present a number of clinical challenges<sup>260</sup>. COPD is a negative prognostic factor for lung cancer patients. The presence of COPD may alter the treatment plan (radiotherapy, surgical approach) for lung cancer. Clinical management of patients with coexistence of COPD and lung cancer requires a multidisciplinary oncology board that includes a pulmonologist. Detailed evaluation (lung function tests, cardiopulmonary exercise test) and management (inhaled drugs, smoking cessation, pulmonary rehabilitation) of COPD should be taken into account for lung cancer treatment (Microablation, surgical approach, palliative therapy, radiotherapy)<sup>261</sup>.

### 11.9 Combined Pulmonary Fibrosis and Emphysema

Combined Pulmonary Fibrosis and Emphysema (CPFE) is suspected based on the presence of upper zone emphysema and lower zone fibrosis<sup>63</sup>




Figure 21: Images of combined pulmonary fibrosis and emphysema

Images from a male smoker aged 77 years with combined pulmonary fibrosis and emphysema demonstrating upper-lobe emphysema and lower-lobe pulmonary fibrosis. A, Chest radiograph. B, CT image. Pulmonary function tests showed an FEV<sub>1</sub> of 108% predicted, FVC of 107% predicted, total lung capacity of 121%, and diffusing capacity of lung for carbon monoxide of 36% predicted. Echocardiography demonstrated grade I diastolic dysfunction and estimated pulmonary artery pressure of 30 to 40 mm Hg<sup>262</sup>.

Spirometry is frequently normal due to opposing effects of hyperinflation and obstruction from emphysema and restriction from fibrosis. Gas transfer however, is usually severely impaired due to the additive effect of dual pathology<sup>263</sup>. Cigarette smoking is a major risk factor for CPFE<sup>264</sup>. It occurs predominantly in males (up to 9:1 male:female ratio). In non-smokers, CPFE has been described in people with occupational dust exposure and genetic mutations<sup>262</sup>.

CPFE is frequently complicated by pulmonary hypertension, lung cancer, acute lung injury and coronary artery disease. Mortality is significant, especially in the presence of pulmonary hypertension<sup>265</sup>.

Even in patients who do not fulfil criteria for IPF, the presence of interstitial features in addition to emphysema carries a significantly higher mortality<sup>63</sup>.

In most cases, high resolution computed tomography (HRCT), spirometry and diffusing capacity of lung for carbon monoxide (DLCO) test are adequate to diagnose CPFE<sup>63</sup>. The prevalence of lung cancer is higher in CPFE than COPD<sup>266</sup>. Therefore, more vigilant follow up of pulmonary nodules is recommended, though no specific screening guideline has been developed for CPFE<sup>262,265</sup>. Squamous Cell Carcinoma was the most common type (42.3%), followed by adenocarcinoma (34.4%).

The median survival for CPFE patients with lung cancer (19.5 months) was significantly shorter than in non-CPFE (53.1 months)<sup>266</sup>.

Currently, no specific treatment exists for CPFE. Post-hoc data from nintedanib trials, which included patients with concurrent emphysema, showed attenuation of rate of decline in forced vital capacity (FVC) in IPF with emphysema, similar to IPF without emphysema<sup>267,268</sup>. An observational cohort study of real-world patients who were commenced on pirfenidone also showed similar rate of progression between CPFE and IPF without emphysema<sup>269</sup>. Hence, anti-fibrotic therapy can be considered in CPFE, where presence of IPF is confirmed, Stem Cell therapy may be a good option for rapidly declining lung function<sup>270</sup>.

# Holistic Management of COPD




Figure 22: Algorithm of COPD MANAGEMENT

## 12 CHAPTER 12: NON-PHARMACOLOGICAL MANAGEMENT

### 12.1 For All Patients of COPD:

Table 18: Non-pharmacological therapy for all COPD

|    |   |    |   |
|----|---|----|---|
| 1. | Asses every patient by MRC dyspnea scale or CAT score and CPET for pulmonary rehabilitation | 2. | Classify COPD with all possible comorbidities and staging |
| 3. | Avoid smoking and smoke   | 4. | Avoid air pollution and temperature: cold and hot         |
| 5. | Immunization  | 6. | Pulmonary Rehabilitation                                  |

For Selected Patients of COPD:

Table 19: Non-pharmacological therapy for specific COPD

|    |  |    |  |
|----|--|----|--|
| 1. | <ul style="list-style-type: none"> <li>• Long Term Oxygen Therapy (LTOT)</li> <li>• Ambulatory Oxygen Therapy</li> </ul> | 2. | Short-Burst Oxygen Therapy   |
| 3. | Non-Invasive Ventilation (NIV):<br>Acute NIV<br>Chronic NIV  | 4. | Surgery:<br>Bullectomy,<br>Lung Volume Reduction Surgery (LVRS) and Endobronchial Procedures |
| 5. | Extracorporeal Membrane Oxygenation  | 6. | Exosome Therapy and Stem cell Therapy  |
| 7. | Lung Transplantation   | 8. | End of Life (EOL) Care:  |

**1. Quit Smoking:** Smoking cessation has the greatest capacity to influence the natural history of COPD

- Nicotine Replacement Therapy (NRT): NRT has a 2-fold increase in smoking cessation rates when compared to placebo. The goal of NRT is to provide the patient with the addictive nicotine without using the harmful tobacco, facilitating avoidance from cigarettes. There are many formulations, including gum, lozenge, transdermal patch, inhaler and a nasal spray. The transdermal patch is the preferred method, as it has the most reliable and steady delivery of nicotine<sup>271</sup>.

## BRIEF STRATEGIES TO HELP THE PATIENT WILLING TO QUIT

- Ask and identify smokers. Document smoking status in the medical record.
- Assess the degree of nicotine dependence and motivation or readiness

### TO QUIT

- Advise smokers about the risks of smoking and benefits of quitting and discuss options
- Assist cessation — this may include specific advice about pharmacological interventions or referral to a formal cessation program.
- Arrange follow-up to reinforce messages.

## 2. Avoid exposure to Smoke or Temperature Fluctuations

**3. Immunization:** Vaccinations are a good way to reduce exacerbations. It has been demonstrated that inactivated influenza vaccines in COPD patients resulted in a significant reduction in the total number of exacerbations per vaccinated subject compared with those who received placebo<sup>272</sup>. Wongsurakiat P et al. described an absolute 21.3% reduction in the incidence of acute respiratory illness following influenza vaccination<sup>273</sup>. Pneumococcal vaccination is recommended to be given in addition to annual influenza vaccination in people with COPD<sup>274</sup>.

## 12.2 Pulmonary Rehabilitation (PR):

### 12.2.1 Patho-physiology of exercise limitation:

Breathlessness on exertion and its phobia is very common in COPD and other Chronic Respiratory Disease (CRD). Inactivity leads to deconditioning, mainly caused by breathlessness which leads to avoidance of physical and social activities, thrusting the patient into a vicious circle leading to further isolation and depression, accompanied by a reduced HRQoL(Fig. 17)<sup>242</sup>. Pathophysiological features of exercise limitation are described in the figure below:




Figure 23: Patho-physiology of exercise limitation  
(Reproduced from Habib GM with permission)

### 12.2.2 Integrative Care in COPD Pulmonary Rehabilitation

Maintaining physical activity and muscle strength is important for every patient suffering from COPD. Physical activity can be achieved through enrollment in an organized program, such as pulmonary rehabilitation (PR). It is a proven effective modality in the treatment of COPD and is part of COPD management guidelines<sup>275</sup>. Ideal candidates for PR are both males and females with moderate, severe or very severe COPD<sup>276</sup>. It should be offered to those patients with COPD who remain symptomatic despite bronchodilator therapy and should be implemented within one month following an acute exacerbation.

- Pulmonary rehabilitation programs involve patient assessment, supervised exercise training, education, behavior change, nutritional intervention and psychosocial support<sup>277</sup>.
- Exercise training is considered to be the cornerstone of pulmonary rehabilitation<sup>277</sup>.
- The benefits of pulmonary rehabilitation include a reduction in symptoms (dyspnea and fatigue), anxiety and depression, and improvements in health-related quality of life (HRQoL), peripheral muscle function and exercise capacity.

- Some patients who experience marked oxygen desaturation on exertion may benefit from ambulatory oxygen during exercise training and activities of daily living
- Duration of pulmonary rehabilitation program in Bangladesh (for example in Pulmo-FIT) ranges from 2 to 8 weeks.
- It is clearly demonstrated that
- Doing any physical activity is better than doing none<sup>63</sup>;
- Be active on most, preferably all, days every week;
- Accumulate 150 to 300 minutes of moderate intensity physical activity or 75 to 150 minutes of vigorous intensity physical activity, or an equivalent combination of both moderate and vigorous activities, each week;
- Do muscle strengthen activities on at least 2 days each week.




Figure 24: COPD's vicious circle (reproduced from<sup>242</sup>).

### 12.2.3 Physical Activity

#### What is physical activity?

Optimal physical activity is the key component to maintaining sustainable benefit of successful PR.

Any bodily movement produced by skeletal muscle that leads to energy expenditure. This includes<sup>278</sup>:

1. At least 30 minutes moderate intensity physical activity for 5 or more days a week.
2. Vigorous physical activities for 20 minutes 3 days every week

3. Shorter bouts three-times moderate 10 minutes or two-times vigorous 10 minutes<sup>278,279</sup>

Physical activity is measured during, work, commuting, and recreation by MET (Metabolic Equivalent Task); sitting 1; moderate 4; vigorous 8. The total PA score is calculated by the sum of all MET/min/week from moderate to severe PA performed in work, commuting, and recreation<sup>280</sup>.

The patterns of PA and energy expenditure differs between the developed and developing countries because of their of their diverse way of life dynamics<sup>281</sup>.

WHO define<sup>282</sup>

**Physical Activity**

“Physical activity” is bodily movement produced by skeletal muscles that requires energy expenditure – including activities undertaken while working, playing, carrying out household chores, travelling, and engaging in recreational pursuits.

**Exercise**

"Exercise" is a subcategory of physical activity that is planned, structured, repetitive, and aims to improve or maintain one or more components of physical fitness.

**Classifying the physical activity level:**

Degree of PA is divided in three groups according to the GPAQ analysis framework: low, moderate, and high<sup>280</sup>.

Table 20: Degree of Physical Activity

| Level of PA |   |
|-------------|---|
| High level  | A person reaching any of the following criteria:<br>(a) Vigorous-intensity activity on at least three days and accumulating at least 1500 MET-minutes/week OR<br>(b) Seven or more days of any combination of walking, moderate- or vigorous-intensity activities accumulating at least 3000 MET-minutes/week |



|          |  |
|----------|--|
| Moderate | A person not meeting the criteria for the 'High' category, but meeting any of the following criteria is classified in this category:<br>1. (a) Three or more days of vigorous-intensity activity of at least 20 min per day OR<br>2. (b) Five or more days of moderate-intensity activity and/or walking of at least 30 min per day OR<br>3. (c) Five or more days of any combination of walking, moderate- or vigorous-intensity activities accumulating at least 600 MET minutes/week. |
| Low      | A person not meeting any of the above-mentioned criteria falls in this category. No activity is reported or some activity is reported but not enough to meet high and moderate categories  |

### Prevalence of PA and Physical inactivity (PiA):

- Globally, 1 in 4 adults is not active enough. 23% of adults aged 18 and over have PiA in 2010 (M-20%, F-27%); in HICs M-26% & F-35%; LMICs M-12%, F-24%.
- More than 80% of world's adolescent population (11-17 years) is insufficiently physically active (M-78%, F-84%)
- Policy for addressing the PiA is taken by 56% of the member countries of WHO
- WHO member countries agreed to reduce PiA by 10% by 2025 PA data from Bangladeshi population<sup>283</sup>:




Figure 25: physical activity in urban and rural of Bangladesh. Composition of total physical activity (%) in urban and rural areas of Bangladesh.

The country wide prevalence of PA levels in a study in Bangladesh (Moniruz-zaman et al 2016) was low 34.5% moderate 46.0% and high 19.5%. Composition of total PA in urban and rural areas<sup>283</sup>:

Table 21:Components of Physical Activity

| Domain                | Rural | Urban |
|-----------------------|-------|-------|
| Workplace activity    | 61%   | 47%   |
| Commuting activity    | 31%   | 38%   |
| Leisure-time activity | 9%    | 15%   |


### Why physical activity is important?<sup>282</sup>

- Regular PA of moderate intensity e.g. walking, cycling, doing sports has significant benefits both in prevention and care of many NCDs.
- It improves skeleto-muscular and cardio-pulmonary fitness
- Improves functional and mental health
- Reduces the risk of hypertension, coronary heart disease, stroke, diabetes, various types of cancer including breast and colon cancer and depression.
- Plays a role both in primary and secondary prevention
- Improves balance impairment and reduce fall incidence
- Optimise balance between energy intake and burning resulting weight control
- Reduces mortality from NCDs

### How to increase PA?

National, local, social, and individual initiative and rationale strategy might increase the PA for people. These might be:

- Awareness and promotion of PA through activity of daily life in association with the relevant sectors of PA
- Planned urbanisation to providing walking, cycling, and other forms of transport accessible and safe for all
- Work place and labour policy should be organised in a way that encourage physical activity
- Sports and PA programme should be organised during free time in school and colleges
- Physical educator should be appointed in every institute
- Sports and recreation facilities involving PA should be arranged to reach as much as people can avail.
- Follow WHO recommendation:
  - o Develop and implement national guideline for PA
  - o Integration of PA with other related policy sectors
  - o Use of mass media to raise awareness of the benefits of PA
  - o Surveillance and monitoring the PA implementation programme



**Assesment of Physical Activity in our setting**

Figure 26: Assessment of Physical Activity in Pulmo-Fit

### 12.2.4 Education and self-management

- Education alone has not been shown to be effective without physical training and self-management plan<sup>284</sup>
- Self-management means intervention with communication with a health care professional to improve health status and decreases hospitalizations and emergency department visits of COPD patients.

## 12.3 Non-Invasive Ventilation (NIV):

### 12.3.1 a. Acute NIV:

Non-Invasive positive pressure ventilation (NIPPV) is a proven modality to decrease morbidity, improve survival and decrease the need for mechanical ventilation in COPD patients who develop acute respiratory failure. Its benefit has been proven in various settings, including the ICU, ward and emergency department in patients with moderate to severe respiratory acidosis<sup>285</sup>. NIV is also effective in facilitating extubation in patients mechanically ventilated for an acute exacerbation. Patients extubated to NIV had fewer re-intubations, fewer tracheostomies, shorter ICU stays, improved ICU survival and fewer complications including nosocomial pneumonia<sup>286</sup>. Because of these positive results, NIV should be considered in patients that require mechanical ventilation for respiratory failure particularly in those who have failed traditional weaning.

### 12.3.2 b. Chronic NIV:

The use of chronic NIV in patients with COPD is more controversial. Struik et al<sup>287</sup> have demonstrated that the use of NIV in COPD improvements in daytime

PaCO<sub>2</sub> and nocturnal transcutaneous PCO<sub>2</sub> measurements in the NIV group. But Number of exacerbations, lung function, mood state, daily activity levels or dyspnoea was not significantly different from standard treatment without COPD. Despite this lack of convincing evidence for its use, there are some clear situations where chronic NIV proves beneficial. One of these situations is in the patient with combination of COPD and obstructive sleep apnea (OSA)<sup>288</sup>. Patients with combined COPD and OSA who are not treated with appropriate NIV, like continuous positive airway pressure (CPAP/BiPAP) for example, experience higher rates of hospitalizations and mortality when admitted with a COPD exacerbation. The use of CPAP/BiPAP for combined OSA and COPD also reduces pulmonary hypertension rates and nocturnal hypoxemia<sup>288</sup>.

#### **12.4 Long Term Oxygen Therapy (LTOT):**

According to the British Thoracic Society (BTS)<sup>289</sup>, long term oxygen therapy is recommended for patients with a PaO<sub>2</sub> ≤55 mm of Hg or ≤7.3kPa or a PaO<sub>2</sub> between 55 mm of Hg or 7.3kPa or 60 mm of Hg ≤8kPa with signs of peripheral edema, polycythemia (hematocrit>55%) or pulmonary hypertension. In this population, long term oxygen therapy is one of the few interventions that prolong life expectancy<sup>290</sup>. It also reduces pulmonary hypertension and hematocrit. Continuous oxygen therapy for 24h/d is superior to nocturnal oxygen in those with severe hypoxemia (PaO<sub>2</sub> ≤7.3kPa or a PaO<sub>2</sub> of ≤8kPa cor-pulmonale)<sup>291</sup>. Situational oxygen is also recommended for those who desaturate with exertion or have hypoxemia at night<sup>292</sup>.

#### **12.5 Ambulatory Oxygen Therapy**

In patients with COPD with moderate hypoxia, current evidence on ambulatory oxygen therapy reveals improvements in dyspnoea post exercise and in the dyspnoea and fatigue domain of quality of life<sup>293</sup>. For patients on LTOT who are motivated to use oxygen outside the home, ambulatory oxygen should be prescribed. Ambulatory oxygen is also indicated for those patients not on LTOT but who have exercise arterial desaturation<sup>292</sup>. In this population, arterial oxygen desaturation is defined as a fall in SaO<sub>2</sub> of 4% to a value <90%. In addition to desaturation, these patients must also demonstrate improvement in exercise capacity and/or dyspnea with oxygen. The goal is to maintain oxygen saturation >90% during exercise.

#### **12.6 Short-Burst Oxygen Therapy**

Short burst oxygen therapy (SBOT) refers to the intermittent use of oxygen, usually from a static cylinder. It is widely prescribed for the alleviation of breathlessness in patients with chronic obstructive pulmonary disease (COPD), despite little convincing evidence of benefit<sup>294</sup>. It should only be considered for episodes of severe breathlessness in patients with COPD not relieved by

inhaler or other reliever treatments. This mode of oxygen delivery should only continue to be prescribed if patient feels improvement in breathlessness with its use<sup>295</sup>.

### 12.7 Extracorporeal Membrane Oxygenation (ECMO):

Extracorporeal membrane oxygenation, or ECMO, is a therapy that adds oxygen to blood and pumps it through their body like the heart. The process takes place outside the body.

ECMO is used when usual treatments are not working. ECMO does not cure heart or lung disease, it only provides time for the patient's heart or lungs to heal<sup>296</sup>.

The ECMO pump pulls blood that has no oxygen attached from a vein and pushes it into the machine's artificial lung, or oxygenator. That's where carbon dioxide is removed from the blood and oxygen is added. There's a color change as the darker blood with no oxygen turns bright red when oxygen is attached to it. As the red blood leaves the oxygenator it is warmed before returning to the patient. Because ECMO is also helping do the work of the patient's lungs, the ECMO team can lower the settings of the ventilator, allowing the lungs to rest and heal.




Figure 27: Mechanism of ECMO

*(Copied from the webpage of the University of Iowa applying for permission) ECMO handles the body's blood in two stages. First it draws blood out of the body and brings it into the ECMO machine for oxygenation and removal of CO<sub>2</sub>. Second it delivers blood from the machine back into the body<sup>296</sup>.*

This is a new modality being used to treat COPD exacerbations to prevent the need for mechanical ventilation and the associated morbidity associated with intubation. As demonstrated above, non-invasive ventilation is superior to mechanical ventilation in terms of mortality, complications and hospital length of stay. However, 15-26% of patients with an acute exacerbation of COPD fails non-invasive ventilation and requires transition to mechanical ventilation. These patients have a higher mortality than those initially treated with mechanical ventilation<sup>276</sup>. There are no current guidelines on the use of ECMO for COPD exacerbations as its use in this indication is new. Studies have shown that the Novalung could prevent intubation in 90% of patients with a trend to decreasing hospital length of stay<sup>289</sup>. It is likely that in the future we will see this modality being used more in centers that have access to it.

## **12.8 Surgery**

None of the current surgical approaches in patients with COPD provides a survival advantage. In view of the potential for serious morbidity and mortality, all surgical treatments require careful assessment by an experienced thoracic medical and surgical team.

### **12.8.1 Bullectomy**

This operation involves resection of large bullae (larger than 5cm). The procedure is most successful where there are very large bulla compressing adjacent apparently normal lung. Giant bullae can be defined as occupying more than 50% of the hemithorax with definite displacement of adjacent lung tissue<sup>297</sup>.

### **12.8.2 Lung Volume Reduction and Endobronchial Procedures:**

LVRS is an effective treatment option in patients with heterogeneous upper zone emphysema and reduced exercise tolerance<sup>298</sup>. According to the National Emphysema Treatment Trial (NETT) trial, LVRS has shown to be superior to medical treatment. LVRS has improved maximal ventilation rate and tidal volume and has lower BORG dyspnea scores<sup>299</sup>. However, surgery carries a high risk of morbidity, particularly in patients with severe COPD, who often have additional co-morbidities. Thus, newer, non-surgical, less invasive strategies have been developed to potentially offer benefit to those patients who would not be candidates for LVRS. These non-surgical procedures are referred to as bronchoscopic lung volume reduction<sup>300</sup>.

### 12.8.3 Lung Transplantation:

Lung transplant is offered to people with untreatable, end-stage lung disease with a limited life expectancy. COPD is one such lung disease and is currently the most common indication for a double lung transplant<sup>301</sup>. According to the International Society for Heart and Lung Transplantation, the disease specific criteria for referral of patients with COPD include an  $FEV_1 \leq 30\%$  while having maximized both pharmacological and non-pharmacological therapy, including bronchodilators, home oxygen, smoking cessation and pulmonary rehabilitation. In addition, they should have a BODE index exceeding 5<sup>302</sup>. Other criteria for transplantation include patients with a BODE index of 7-10 plus a history of hospitalization for exacerbation associated with acute hypercapnia, pulmonary hypertension or cor pulmonale or an  $FEV_1$  of less than 20% with either a DLCO of less than 20% or a homogeneous distribution of emphysema<sup>301</sup>.

The survival benefit for patients with COPD receiving a transplant is not as robust as for transplant in other respiratory conditions, such as pulmonary fibrosis (DPLD), cystic fibrosis and primary pulmonary hypertension<sup>302</sup>. If a patient seems like an appropriate candidate and is willing, a referral to a transplant centre should be made.

### 12.9 End of Life (EOL)Care:

Patients with COPD typically experience the last few months to years of their life marked by progressive dyspnea, functional decline and social isolation<sup>302</sup>. Their quality of life has been compared to those with lung cancer<sup>303</sup>. As with any non-curable chronic illness, goals of care and end of life discussions must occur<sup>304</sup>. The discussion should focus on education of the disease course and management of end of life issues, such as symptom control and advanced health care directives. The prognosis of COPD is often difficult to predict as the disease is marked by progressive decline in lung function with increasing symptoms. There are episodes of acute illness and associated co-morbid conditions. Because prognosis is difficult to predict, it is often unclear as to when to begin end of life care discussions and palliative symptom management<sup>305</sup>. These discussions should definitely occur when the  $FEV_1$  is  $<30\%$ , abnormal blood gases are present or when cor-pulmonale with pulmonary hypertension develops as these patients are found to have the poorest prognosis. The major symptoms experienced by end-stage COPD patients include dyspnea, pain, fatigue and insomnia<sup>305</sup>. Insomnia is often improved simply through dyspnea control<sup>305</sup>. Low dose Benzodiazepines can be used as adjuncts to opioids to control severe dyspnea along with BiPAP and oxygen.

### 13.1 Pharmacotherapy in COPD

The current pharmacological treatment of COPD is symptomatic and is mainly based on bronchodilators, such as selective Beta<sub>2</sub>-adrenergic agonists (short- and long-acting), anticholinergics, theophylline, phosphodiesterase-4 antagonists, such as roflumilast or a combination of these drugs<sup>306</sup>. Inhaled corticosteroids (ICS) are not generally recommended for patients with COPD without features of Asthma due to their lack of efficacy, side effects, and high costs. However, oral or parenteral glucocorticoids may be given during exacerbations of COPD. A number of novel long-acting anticholinergics e.g. Glycopyrronium and Beta<sub>2</sub>-agonists e.g. Indacaterol/with once daily are now clinically using<sup>307</sup>. Montelukast has been not yet been established in chronic obstructive pulmonary disease (COPD)<sup>308</sup>

Patients with characteristics of both COPD and asthma, sometimes referred to as asthma COPD overlap (ACO), in our classification COPD -A may be more likely to benefit from montelukast therapy<sup>309</sup>

Table 22: Pharmacotherapy in COPD

|                                    |   |
|------------------------------------|---|
| Stage 1:<br>FEV <sub>1</sub> ≥ 80% | SMART THERAPY<br>As per need Budesonide and Formoterol inhaler therapy <sup>310</sup>   |
| Stage 2<br>FEV <sub>1</sub> <80-50 | TRIPPLE THERAPY <sup>311</sup><br>(LAMA+LA-BA+ICS) (Moderate Dose)<br>(FEV <sub>1</sub> <80% to ≥ 60% to Maximum dose if FEV <sub>1</sub> ≤60%)<br>and + add on ± Montelukast to control COPD A plus, Ipratropium & salbutamol as per need <sup>312</sup> |
| Stage 3<br>FEV <sub>1</sub> <50-30 | TRIPPLE THERAPY <sup>311</sup><br>(LAMA+LABA+ICS)(Maximum Dose) +Doxophylline <sup>313</sup><br>± Mon-telukast <sup>309</sup> plus Ipratropium & salbutamol as per need <sup>312</sup>  |
| Stage 4<br>FEV <sub>1</sub> <30    | TRIPPLE THERAPY <sup>311</sup><br>(LAMA+LA-BA+ICS)(Maximum Dose)+Mon-telukast + Doxophylline <sup>313</sup> + Nebulized Bronchodilator + Neb Budesonide <sup>314, ± LTOT<sup>294</sup></sup>  |



Table 23: Pharmacotherapy for COPD-B

|  |
|--|
| <p><b>Stage:1</b><br/>As per need Salbutamol and ipratropium inhalation<sup>311</sup></p>  |
| <p><b>Stage:2</b><br/>DOUBLE THERAPY:<sup>307</sup><br/>(LAMA+LABA) +Sub-lingual Lyophilized Bacterial Lysate<sup>138</sup> (10 days / months &amp; 3 months/ year)</p>  |
| <p><b>Stage:3</b><br/>DOUBLE THERAPY:<br/>(LAMA+LABA)+Doxophylline<sup>313</sup> +Sublingual Lyophilized Bacterial Lysate (10 days /month &amp; 3 months/ year) +Azithromycin<sup>140</sup> (alternate day or thrice a week throughout the year)</p>   |
| <p><b>Stage:4</b><br/>DOUBLE THERAPY:<br/>(LAMA+LABA) +Roflumilast<sup>306</sup>+Sublingual Lyophilized Bacterial Lysate (10 days /month &amp; 3 months/ year) +Azithromycin (alternate day or thrice a week throughout the year) +Tobramycin<sup>135</sup> nebulization 3 weeks 12 hourly then 3 weeks gap life-long ± LTOT<sup>289</sup></p> |

Table 24:Pharmacotherapy of COPD-G

|  |
|--|
| <p>Stage:1 PPI<sup>149,150</sup> + Domperidone<br/>C h a n g e food habit Sleep with double pillow Eating &amp; drinking slowly to avoid cough, wheeze or choking plus As per need Salbutamol and ipratropium inhalation<sup>311</sup></p>   |
| <p>Stage:2 DOUBLE THERAPY: (LAMA+LA-BA)+PPI +DOMPERIDON Change food habit Sleep with double pillow Eating &amp; drinking slowly to avoid cough, wheeze or choking. Plus, as per need Salbutamol and ipratropium inhalation<sup>311</sup></p>   |
| <p>Stage:3 Avoid theophylline / Doxophylline as it may cause GERD<sup>315</sup><br/>DOUBLE THERAPY:<br/>(LAMA+LABA) +Nebulize Bronchodilator 1-2 times daily +PPI +DOMPERIDON Change food habit Sleep with double pillow Eating &amp; drinking slowly to avoid cough, wheeze or choking. + as per need Salbutamol and ipratropium inhalation<sup>311</sup></p> |
| <p>Stage:4 DOUBLE THERAPY: (LAMA+LABA)+Nebulize Bronchodilator 3-4 times daily + PPI +DOMPERIDON Change food habit Sleep with double pillow Eating &amp; drinking slowly to avoid cough, wheeze or choking . ± LTOT</p>  |

Table 25: Pharmacotherapy of COPD - O & S

| Stages  | COPD – O   | COPD - S   |
|---|--|--|
| <b>Stage 1:</b><br><b>FEV<sub>1</sub> &gt;80%</b> | Weight Reduction and CPAP <sup>170,171</sup> plus As per need Salbutamol and ipratropium inhalation <sup>311</sup>           | As per need Salbutamol and ipratropium inhalation <sup>311</sup>                                       |
| <b>Stage 2:</b><br><b>FEV<sub>1</sub> 50-79%</b>  | <b>Double therapy:</b><br>(LAMA+LABA) Always sleep in lateral Position<br>-Add CPAP or BiPAP plus As per need Salbutamol and | <b>Double therapy:</b><br>(LAMA+LABA)  |
| <b>Stage 3:</b><br><b>FEV<sub>1</sub> 30-49%</b>  | Double therapy:<br>(LAMA+LABA) +Doxophylline<br>Always sleep in lateral Position ± add CPAP or BiPAP                         | <b>Double therapy:</b><br>(LAMA+LABA) +Doxo-phylline<br>Always sleep in lateral Position ± add CPAP or |
| <b>Stage 4:</b><br><b>FEV<sub>1</sub> &lt;30%</b> | Double therapy: (LAMA+LA-BA)+Doxophylline +Nebulize Bronchodilator<br>Always sleep in lateral Position ± add BiPAP ± LTOT    | Double therapy:<br>(LAMA+LABA)<br>+ Doxophylline+Nebulize Bronchodilator ± LTOT                        |

Table 26: Pharmacotherapy of COPD - U

| COPD – U                                    |   |
|---|---|
| <b>Stages</b>                               | Practically nil case stage 1 with Unstable COPD   |
| <b>Stage 1:<br/>FEV<sub>1</sub> &gt;80%</b> | TRIPPLE THERAPY<br>(LAMA+LA-BA+ICS) (Maximum Dose)  |
| <b>Stage 2:<br/>FEV<sub>1</sub> 50-79%</b>  | TRIPPLE THERAPY<br>(LAMA+LA-BA+ICS) (Maximum Dose)<br>+Montelukast+Doxophylline<br>+Azithromycin (alternate day or thrice a week throughout the year)<br>+Sublingual Lyophilized Bacterial Lysate (10 days /month & 3 months/ year) |
| <b>Stage 3:<br/>FEV<sub>1</sub> 30-49%</b>  | COMBINE ALL (for maximum relief of Symptoms) + LTOT + BIPAP during sleep  |
| <b>Stage 4:<br/>FEV<sub>1</sub> &lt;30%</b> |   |

### 13.2 Treatment of anxiety and depression in COPD

**Anxiety and depression** are the most **common** psychiatric **comorbidities** associated with **COPD**. Varying prevalence rates have been reported from various parts of world ranging from 5% to more than 40% both because of variation in the study participants to various psychological tools having been used for screening<sup>316,317</sup>.

Anxiety symptoms in COPD are associated with worse quality of life<sup>318</sup> selfmanagement and exercise performance and with increased medical symptom reporting<sup>63</sup> exacerbations hospitalizations, length of hospitalizations, medical costs and mortality<sup>318</sup>. The prevalence of one anxiety disorder in particular, is panic disorder, it is approximately 10 times greater in COPD than the population prevalence of 1.5 to 3.5%, and panic attacks are commonly

experienced<sup>319</sup>.

Selective serotonin reuptake inhibitors (SSRIs) such as sertraline, have been recommended as first line pharmacological therapies for anxiety in COPD<sup>320</sup>.

**Benzodiazepine** users **were** at “significantly **higher risk**” for outpatient respiratory **exacerbations**, emergency room visits for either **COPD** or pneumonia, and that hospitalizations among this group compared to nonusers<sup>321</sup>.

Table 27: Classification of Benzodiazepine<sup>322</sup>

|                       | Physical Findings | Investigation   |
|-----------------------|-------------------|---|
| Half-life designation | Half-life (h)     | Name of Drugs   |
| Long-acting           | 25–100            | Bromazepam, chlordiazepoxide, clobazam, clonazepam, clorazepate, diazepam, flurazepam, nitrazepam |
| Intermediate-acting   | 5–24              | Alprazolam, lorazepam, oxazepam, temazepam  |
| Short-acting          | < 5               | Triazolam, Midazolam  |

Among individuals with more severe COPD, 35.4 % of incident (Aspiration, Pneumonia, Respiratory Failure) use occurred during a COPD exacerbation<sup>322</sup>. In severe COPD hospitalizations due to exacerbations and pneumonia risks were “significantly elevated” in COPD patients that used benzodiazepines.

The potential adverse respiratory effects of benzodiazepines in COPD may also be heightened in older adults given their altered pharmacokinetics that increase benzodiazepine half-life<sup>322</sup>.

Adverse effects associated with benzodiazepine use in COPD patients, may include hypoxemia, decreased respiratory drive, hypercapnea, and decreased respiratory muscle strength<sup>323</sup>.

**NOTE: Low dose benzodiazepines and opioids with Oxygen may be safe for severe COPD and terminally ill patient. But at higher doses it might increase mortality<sup>324</sup>.**

## 14. CHAPTER 14: MANAGEMENT OF EXACERBATIONS OF COPD (Plan ABCO)

Acute exacerbations of chronic obstructive pulmonary disease (COPD) are treated with oxygen (in hypoxemic patients), inhaled beta<sub>2</sub> agonists, inhaled anticholinergics, antibiotics and systemic corticosteroids. Methylxanthine therapy may be considered in patients who do not respond to other bronchodilators<sup>325</sup>. Antibiotic therapy is directed at the most common pathogens, including *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Moraxella catarrhalis*. Mild to moderate exacerbations of COPD are usually treated with older broad-spectrum antibiotics such as amoxicillin-clavulanate potassium, doxycycline, and Levofloxacin. Treatment with augmented penicillins, fluoroquinolones, third-generation cephalosporins and/ or aminoglycosides may be considered in patients with more severe exacerbations and/or *Pseudomonas* infection<sup>326,327</sup>.

### 14.1 Exacerbations of COPD<sup>3</sup>

An exacerbation of chronic obstructive pulmonary disease (COPD) is defined as an acute worsening of respiratory symptoms that results in additional therapy. COPD exacerbations are complex events usually associated with **increased airway inflammation** presented as increased cough and wheeze, **increased mucus production** presented as excess sputum production and sputum color changes and **marked gas trapping in lung** presented as markedly increased dyspnea. As other comorbidities that may worsen respiratory symptoms are common in COPD patients, clinical assessment to rule out differential diagnoses should be considered before diagnosis of a COPD exacerbation.

Always exclude following conditions before establishing diagnosis of acute exacerbations of COPD

**These are 5 P and 1 C: 5 P:** 1. Pneumonia, 2. Pneumothorax, 3. Pleural effusion, 4. Pulmonary embolism, 5. Pulmonary edema and **1.C:** Cardiac arrhythmias.  
Differential diagnosis of acute exacerbations of COPD:

## P1: Pneumonia

### History and Symptoms

Table 28: Diagnostic features of Pneumonia

|   |                           |   |
|---|---------------------------|---|
| History of running nose or sore throat          | Temp more than 99°F       | Chest Xray<br>CBC   |
| Fever, shortness of breath and increasing cough | Tachypnea<br>Tachycardia  | Sputum for C/S  |
| Dull Chest pain                                 | May be Low pressure       | C Reactive protein  |
| Scanty Hemoptysis                               | Features of consolidation | Procalcitonin high in bacterial pneumonia. (optional) If suspect, viral or atypical pneumonia |

## P2: Pneumothorax

Table 29: Diagnostic features of Pneumothorax

| History and Symptoms  | Physical Findings                    | Investigations                  |
|---|--------------------------------------|---------------------------------|
| Sudden onset sharp, one-sided chest pain  | Tachypnea<br>Tachycardia<br>Cyanosis | Chest X-ray<br>Chest ultrasound |
| Sudden shortness of breath  | May be Low pressure                  | CT scan Chest                   |
| <b>Percussion:</b> hyperresonance on one side of the chest. It is like Tympanic sound: hollow, high, drum like sounds |                                      |                                 |
| <b>Auscultation:</b> Breath sound diminished or absent  |                                      |                                 |

**Pneumothorax in COPD:** Spontaneous pneumothorax is defined by the presence of air in the pleural cavity without history of trauma. This is a significant clinical problem. COPD is a common cause of pneumothorax. The risk of recurrence of spontaneous pneumothorax secondary to COPD is high and various studies quote rates 20-60%.

### P3: Pleural Effusion;

Table 30: Diagnostic features of Pleural effusion

|  |                               |  |
|--|-------------------------------|--|
| COPD with SOB usually admitted in hospital or in ICU for SOB | Unilateral<br>Or<br>Bilateral | Chest X-ray<br>Chest ultrasound<br>CT scan Chest<br>Echocardiography |
| Most are due to CHF and/or parapneumonic effusions           |                               | Transudate – CHF Exudate – parapneumonic                             |
| With or without fever  |                               | BNP or Pro-BNP   |

### P4: Pulmonary Embolism

Table 31: Features of Pulmonary embolism

| History and Symptoms                    | Physical Findings  | Investigations   |
|---|--|--|
| Usually patient of COPD with sudden SOB | No change in physical findings except occasional P2 Loud | D-Dimer<br>CT scan with contrast<br>Duplex study of lower extremities<br>CT Pulmonary Angiogram (Optional) |

### P5: Pulmonary edema

Table 32: Features of Pulmonary edema

| History and Symptoms   | Physical Findings   | Investigations   |
|--|---|--|
| Disproportionate shortness of breath and marked limitation of activities despite taking adequate Dual bronchodilator LABA/LAMA | High or Low pressure<br>Tachycardia<br>Arrhythmias like Atrial Fibrillation             | <ul style="list-style-type: none"> <li>• X-ray Chest</li> <li>• USG of both chest</li> <li>• BNP or NT-pro BNP</li> <li>• Colour doppler Echocardiogram</li> </ul> |
| History of old MI/ Angina like chest pain during exertion/ PND/ Orthopnea  | Basal Crepts and ronchi with prolong expiration associated features of Pleural Effusion |  |

## C1: Cardiac Arrhythmias

Table 33: Identifying Cardiac arrhythmias

|  |  |   |
|--|--|---|
| Feelings of palpitation<br>Sudden feelings of extra beats with SOB | Irregular pulse e.g. Atrial Fibrillation | <ul style="list-style-type: none"> <li>• ECG</li> <li>• Telecardiac monitoring during 6 minute exercise test/ Incremental Schuttle test or Cardio-Pulmonary Exercise Tests</li> </ul> |
|--|--|---|

### 14.2 COPD Exacerbations PLAN (ABCO):

Antibiotics (A), Bronchodilator (B), Systemic corticosteroids (C) and Oxygen (O): ABCO

### 14.3 A. Antibiotics for treatment of Exacerbations:

Exacerbations with clinical features of infection (increased volume and change in color of sputum and/or fever) usually will benefit from antibiotic therapy. Bacterial infection may have either a primary or secondary role in about 50% of exacerbations of COPD<sup>328,329</sup>. Haemophilus influenzae, Streptococcus pneumoniae and Moraxella catarrhalis are most commonly involved<sup>330</sup>. Mycoplasma pneumoniae and Chlamydia pneumoniae have also been reported<sup>328,331</sup>. As lung function deteriorates (FEV<sub>1</sub><35%), Pseudomonas aeruginosa and Staphylococcus aureus are often encountered<sup>326,332</sup>. Multi drug resistant Ps. aeruginosa is associated with 6 fold increased risk of death<sup>333</sup>. It is clearly demonstrated that COPD exacerbations treated in primary care, use of point-of-care C-Reactive Protein (CRP) testing to guide prescribing of antibiotics lowered patient- reported antibiotic use (OR 0.31, 95% CI 0.20 to 0.47)<sup>334</sup>. The judicious use of CRP testing in primary or tertiary care may assist in determining the need for antibiotics for exacerbation management.

Table 34<sup>335</sup>: Cut off Values of CRP for COPD needs antibiotics

|  |  |
|--|--|
| Typical clinical cut off concentrations: Conventional C Reactive Protein (CRP) | Cutoff: approximately 10 mg/L. Apparently healthy individuals: less than or equal to 5 mg/L Acute range: 20-500 mg/L |
|--|--|

**Procalcitonin (PCT)** is an acute phase reactant. Procalcitonin levels increase in bacterial infections but do not increase in viral infections or auto-immune inflammation<sup>336</sup>. Procalcitonin has been proposed as a measure to determine if



patients with an exacerbation of COPD require oral antibiotics. In most clinical trials, use of antibiotics was discouraged if procalcitonin was 0.1ng/ml or lower and encouraged if procalcitonin was above 0.25ng/ml (Figure 19)<sup>337</sup>.




Figure 28: PCT in different infections

Available evidence concerning PCT in different infections derived from observational and randomized-controlled intervention studies. While for some infections, intervention studies have investigated benefit and harm of using PCT for antibiotic decisions (right side), for other infections only results from diagnostic (observation) studies are available with mixed results (left side). Abbreviations: PCT, procalcitonin. + moderate evidence in favor of PCT; ++ good evidence in favor of PCT; +++ strong evidence in favor of PCT; ? evidence in favor or against the use of PCT still undefined<sup>337</sup>.

## PCT algorithm for patients with respiratory tract infection





Figure 29: PCT algorithm with respiratory tract infections in A & E  
PCT algorithm in patients with respiratory tract infections in the Emergency Department. The clinical algorithm for antibiotic stewardship in patients with respiratory tract infections in the Emergency Department encourages ( $>0.5 \mu\text{g/l}$  or  $>0.25 \mu\text{g/l}$ ) or discourages ( $<0.1 \mu\text{g/l}$  or  $<0.25 \mu\text{g/l}$ ) initiation or continuation of antibiotic therapy more or less based on PCT specific cutoff ranges. Abbreviations: AB, antibiotic; LRTI, lower respiratory tract infection; PCT, procalcitonin; PSI, Pneumonia Severity Score<sup>337</sup>.

## PCT algorithm for stopping antibiotics in patients with sepsis in the ICU




Figure 30: PCT algorithm with sepsis in the ICU

PCT algorithm in patients with sepsis in the ICU. In critically ill patients in the ICU, cut-offs are higher and initial empiric antibiotic therapy should be encouraged in all patients with suspicion of sepsis. PCT cut-offs are helpful in the subsequent days after admission to shorten the courses of antibiotic therapy in patients with clinical improvement. Abbreviations: AB, antibiotic; PCT, procalcitonin<sup>337</sup>.

## 14.4 systemic corticosteroids + antibiotics for exacerbation

A randomized placebo-controlled trial has provided evidence to support the traditional practice of treating exacerbations with a combination of systemic corticosteroids and antibiotics<sup>338</sup>. In this study, hospitalised patients were commenced on a tapering dose of prednisolone and randomized to receive doxycycline 200mg daily or placebo for 7 days. Clinical cure, defined as complete resolution of signs and symptoms, at day 10 was significantly higher in the antibiotic treated group compared to placebo (OR 1.9, 95%CI 1.2 to 3.2, NNT = 7, 95% CI 4 to 523)<sup>338</sup>.

Therapeutic guidelines: antibiotic recommend the use of oral agents such as Co-amoxycylav or Levofloxacin or Doxycycline for volume and/or color change of sputum in COPD with CRP 20 or more<sup>335</sup>. If patients do not respond to the above antibiotics or If pneumonia, Acinetobacter, Pseudomonas or staphylococci are suspected, appropriate antibiotics should be used<sup>339,340</sup>.

## 14.5 B. Inhaled bronchodilators in exacerbations

### 14.5.1 Effectiveness of Inhaled BD COPD exacerbations

In exacerbations of COPD, the immediate bronchodilator effect is small, but may result in significant improvement in clinical symptoms in patients with severe obstruction.

Studies of acute airflow limitation in asthma indicate that beta-agonists are as effectively delivered by metered dose inhaler and spacer as by nebulizer<sup>341</sup> but in COPD this effect is unknown.

People with COPD often have cardiac co-morbidities, although these may be undiagnosed at the time of presentation with a COPD exacerbation. Such patients may be susceptible to adverse events from high dose, frequent short acting beta agonists. A systematic review demonstrated that higher (5mg versus 2.5mg) doses of salbutamol were associated with increased risk of tremors, elevated heart rate, palpitations and lower blood pressure, but without evidence of any additional benefit of bronchodilatations<sup>342</sup>.

It is advised to use 2.5mg sabutamol nebulization at a time to avoid cardiac toxicity.

In Bangladesh Combination (Ipratropium 500µg and Salbutamol 2.5 mg/3ml) nebulizer solution is used to control symptoms of exacerbations<sup>343</sup>.

### 14.5.2 C. Short course Systemic corticosteroids

Walters et al reported that there is high-quality evidence that systemic corticosteroids reduce (around 5 days) the severity of exacerbations and shorten duration of recovery from exacerbations<sup>344</sup>.

1. Treatment failure (defined as additional treatment, hospital admission/re-admission for index episode, return to emergency department, unscheduled physician visit for the index episode),
2. Improve lung function,
3. Shorten duration of recovery
4. And reduce the severity of exacerbations of COPD. In summary, a 5-day course of oral prednisolone of 30mg to 40mg is adequate to control COPD exacerbations. In patients who have been on oral corticosteroids for longer than 14 days, tapering may be necessary. Patients on long-term oral corticosteroid therapy (>7.5 mg prednisolone daily for more than 6 months) are at risk of developing osteoporosis. Prevention and treatment of corticosteroid-induced osteoporosis should always be considered.

## **14.6 O. Oxygen during exacerbations:**

### **14.6.1 Controlled oxygen therapy:**

This is indicated in patients with hypoxia, with the aim of improving oxygen saturation ( $SpO_2$ ) to 88 to 92%. Use nasal prongs at 1–2.0 L/minute or a face mask at 2 to 4 L/minute. Minimize excessive oxygen administration, which can worsen hypercapnia.

### **14.7 Non-Invasive Ventilatory assistance:**

**BiPAP** is indicated for treating hypercapnia and acidosis. This Non-invasive ventilation by means of a mask is the preferred method.

### 15.1 Pre-operative Assessment for COPD patient:

Patients with COPD are at increased risk of post-operative pulmonary complications after any thoracic or non-thoracic surgery. COPD was independently associated with higher postoperative length of hospital stay, morbidity and mortality<sup>345</sup>, including abdominal operations<sup>346</sup>. Careful pre-operative work-up of patients with COPD minimises post-operative complications.

#### 15.1.1 Lung function assessment for lung surgery

In all patients on the basis of measured preoperative lung functional parameters, we predicted postoperative lung function<sup>347,348</sup>.

In all patients the residual pulmonary function was predicted as follows: it is assumed that each segment of the lung, and there are total 19 of them (upper right lobe - 3, right middle lobe - 2, right lower lobe - 5; upper left lobe - 3, lingula - 2, lower left lobe - 4 segments) in the healthy lung contributes equally to ventilation that is 1/19 (nineteenth part) or 5.26% of total lung ventilation (100%).

Predicted postoperative FEV<sub>1</sub> (Ppo FEV<sub>1</sub>) in absolute values (1-L) is calculated as follows:

- $Ppo\ FEV_1 = \text{preoperative } FEV_1 \times (19 \text{ segments} - \text{the number of segments to be removed}) \div 19$  or by the formula.
- $Ppo\ FEV_1 = \text{preoperative } FEV_1 \times (1 - (S \times 5.26)) \div 100$ . (S = number of segments to be removed).

Both formulas provide the same result.

Ppo FEV<sub>1</sub> should be 1L or more for Preoperative fitness

Other tests:

If performance on the low technology exercise test is satisfactory (stair climbing altitude > 22 m or shuttle walk distance > 400 m), patients are regarded as at low risk of anatomic resection.

**16.1      A. Stem cell therapy for (COPD)**

Stem cells have the unique ability to change into any cell in the body, called differentiation. Studies have shown that expanded cord tissue-derived mesenchymal cells have incredibly high anti-inflammatory properties and tissue regeneration capabilities. When delivered, intravenously MSCs (mesenchymal stem cells) will travel to the lungs (or any area of inflammation) and combat the issue. They can be administered intravenously and find damaged tissues around organs. When used in regards to COPD patients, stem cells can repair damaged lung tissue to combat emphysema or chronic bronchitis. Stem cells have natural anti-inflammatory properties, which can clear airways for those with chronic bronchitis. Completed studies have shown the ability to quantify the effects of stem cell therapy. According to a report done by the Lung Institute called Autologous Stem Cell Therapy and its Effects on COPD, over 82% of patients that attempted stem cell treatment had noticeable improvements in their quality of life after their therapies. Many of these patients reported increased lung capacity and the ability to walk following transfusion. These reports have positioned stem cells as one of the best viable options for current patients with COPD. While stem cell treatment has not shown the ability to cure COPD, its ability to repair damaged tissue and relatively invasive nature makes it an attractive alternative to patients. This path becomes even more desirable when considering the current treatment protocols requiring the use of ventilators or even total lung transplants, which are rare and extremely expensive. But no published data yet recommended for its general use for COPD<sup>349</sup>.

**16.2      B. Exosome in COPD :**

Stem cells-based therapy may be a prospective way for diseases that are irreversible and incurable at present<sup>350</sup>. Specifically, regenerative medicine contains two goals: one is efficiently and safely transferring stem cells into injured organs and tissues, which may replace the transplantation of the entire organ in the near future; the other is to develop strategies in order to improve the regenerative potential and function of adult stem cells residing in various organs<sup>351</sup>.

A key component of paracrine secretion is extracellular vesicles (EVs), particularly the exosome fraction that mainly contributes to the action of stem cells in which genetic information can be horizontally transferred between stem cells and tissue-injured cells. On the basis of the ability of microvesicles (MVs) to mimic stem cell properties, it is speculated that stem cell-derived MVs especially exosomes represent a relevant therapeutic option in regenerative medicine<sup>352</sup>. But still need further effort to explore exact therapeutic use of exosome for COPD<sup>350</sup>.

## 17 CHAPTER 17. References:

- 1 Hassan1, M. R. SEF Staging: A Novel Approach to Optimal COPD Management. *EC PULMONOLOGY AND RESPIRATORY MEDICINE* 2.5 163-171 (2016).
- 2 Cavaillès, A. et al. Comorbidities of COPD. *European Respiratory Review* 22, 454-475 (2013).
- 3 GOLD. Global Initiative for Chronic Obstructive Lung Disease 2020 Report. (2020).
- 4 WHO. Burden of COPD. <<https://www.who.int/respiratory/copd/burden/en/>> (2020).
- 5 Collaborators, G. C. R. D. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet. Respiratory Medicine* 5, 691 (2017).
- 6 WHO. Chronic obstructive pulmonary disease (COPD), <[https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-\(copd\)](https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-(copd))> (2017).
- 7 WHO. Chronic obstructive pulmonary disease (COPD), <[https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-\(copd\)](https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-(copd))> (2017).
- 8 Hassan, M. in *PULMOCON 2008*.
- 9 Perng, D.-W. & Chen, P.-K. The relationship between airway inflammation and exacerbation in chronic obstructive pulmonary disease. *Tuberculosis and respiratory diseases* 80, 325-335 (2017).
- 10 Vestbo, J. & Hansen, E. F. Airway hyperresponsiveness and COPD mortality. *Thorax* 56, ii11-ii14 (2001).
- 11 Martínez-García, M.-A. et al. Prognostic value of bronchiectasis in patients with moderate-to-severe chronic obstructive pulmonary disease. *American journal of respiratory and critical care medicine* 187, 823-831 (2013).
- 12 Liang, B., Wang, M., Yi, Q. & Feng, Y. Association of gastroesophageal reflux disease risk with exacerbations of chronic obstructive pulmonary disease. *Diseases of the Esophagus* 26, 557-560 (2013).
- 13 Marin, J. M., Soriano, J. B., Carrizo, S. J., Boldova, A. & Celli, B. R. Outcomes in patients with chronic obstructive pulmonary disease and obstructive sleep apnea: the overlap syndrome. *American journal of respiratory and critical care medicine* 182, 325-331 (2010).
- 14 al, H. M. e. in *PULMOCON 2008*.
- 15 RA., H. M. a. K. “Newer Classification of COPD and Management Strategy”. . *EC Pulmonology and Respiratory Medicine* 9.2, 01-15 (2020).
- 16 VK, V. Chronic obstructive pulmonary disease. *Indian J Med Res.* 2, 251–269. (2013).
- 17 H, F. J. Emphysema Medical disorder,, <<https://www.britannica.com/science/emphysema>> (2018).
- 18 Marsh, S. E. et al. Proportional classifications of COPD phenotypes. *Thorax* 63, 761-767, doi:10.1136/thx.2007.089193 (2008).
- 19 Taylor, D. R. & Murray, S. A. Improving quality of care for end-stage respiratory disease: changes in attitude, changes in service. *Chronic respiratory disease* 15, 19-25 (2018).
- 20 Stewart, J. I. & Criner, G. J. The small airways in chronic obstructive pulmonary disease: pathology and effects on disease progression and survival. *Current opinion in pulmonary medicine* 19, 109-115 (2013).
- 21 Hogg, J. C., Paré, P. D. & Hackett, T.-L. The contribution of small airway obstruction to the pathogenesis of chronic obstructive pulmonary disease. *Physiological reviews* 97, 529-552 (2017).
- 22 Johns, D. P., Walters, J. A. & Walters, E. H. Diagnosis and early detection of COPD using spirometry. *Journal of thoracic disease* 6, 1557 (2014).
- 23 Sodemán, W. S. T. *Instructions for Geriatric Patients*. Third edn, 464 (2005).



- 24 Odeyemi, Y. E., Ngwa, J., Dodd, K., Gillum, R. F. & Mehari, A. Does Low FEV<sub>1</sub> in Addition to Fixed Ratio and/or Lower Limit of Normal of FEV<sub>1</sub>/FVC Improve Prediction of Mortality in COPD? The NHANES-III-linked-mortality Cohort. *Journal of the National Medical Association* 111, 94-100 (2019).
- 25 Rodriguez-Roisin, R. et al. Chronic respiratory symptoms with normal spirometry. A reliable clinical entity? *American Journal of Respiratory and Critical Care Medicine* 195, 17-22 (2017).
- 26 S, W. in *Medical News Today*, (2018).
- 27 Salvi, S. S. & Barnes, P. J. Chronic obstructive pulmonary disease in non-smokers. *Lancet* 374, 733-743, doi:10.1016/s0140-6736(09)61303-9 (2009).
- 28 Zhang, J., Lin, X.-f. & Bai, C.-x. Comparison of clinical features between non-smokers with COPD and smokers with COPD: a retrospective observational study. *International journal of chronic obstructive pulmonary disease* 9, 57 (2014).
- 29 Restrepo, M. I., Sibila, O. & Anzueto, A. Pneumonia in patients with chronic obstructive pulmonary disease. *Tuberculosis and respiratory diseases* 81, 187-197 (2018).
- 30 Biswas, R. S. R. et al. Indoor biomass fuel smoke exposure as a risk factor for chronic obstructive pulmonary disease (COPD) for women of rural Bangladesh. *Chattagram Maa-O-Shishu Hospital Medical College Journal* 15, 8-11 (2016).
- 31 Blanc, P. D. et al. Occupational exposures and the risk of COPD: dusty trades revisited. *Thorax* 64, 6-12 (2009).
- 32 Diaz-Guzman, E., Aryal, S. & Mannino, D. M. Occupational chronic obstructive pulmonary disease: an update. *Clinics in chest medicine* 33, 625-636 (2012).
- 33 Douwes, J., Thorne, P., Pearce, N. & Heederik, D. Bioaerosol health effects and exposure assessment: progress and prospects. *The Annals of occupational hygiene* 47, 187-200 (2003).
- 34 Li, T. et al. Fine particulate matter (PM<sub>2.5</sub>): The culprit for chronic lung diseases in China. *Chronic diseases and translational medicine* 4, 176-186 (2018).
- 35 Zhu, R. et al. The relationship between particulate matter (PM<sub>10</sub>) and hospitalizations and mortality of chronic obstructive pulmonary disease: a meta-analysis. *COPD: Journal of Chronic Obstructive Pulmonary Disease* 10, 307-315 (2013).
- 36 Hart, J. E., Eisen, E. A. & Laden, F. Occupational diesel exhaust exposure as a risk factor for COPD. *Current opinion in pulmonary medicine* 18, 151 (2012).
- 37 Mehta, A. J. et al. Occupational exposure to dusts, gases, and fumes and incidence of chronic obstructive pulmonary disease in the Swiss Cohort Study on Air Pollution and Lung and Heart Diseases in Adults. *American journal of respiratory and critical care medicine* 185, 1292-1300 (2012).
- 38 Molfino, N. A. & Coyle, A. J. Gene-environment interactions in chronic obstructive pulmonary disease. *International journal of chronic obstructive pulmonary disease* 3, 491 (2008).
- 39 Anthonisen, N. R. et al. The effects of a smoking cessation intervention on 14.5-year mortality: a randomized clinical trial. *Annals of internal medicine* 142, 233-239 (2005).
- 40 Molfino, N. A. & Coyle, A. J. Gene-environment interactions in chronic obstructive pulmonary disease. *International journal of chronic obstructive pulmonary disease* 3, 491-497, doi:10.2147/copd.s2528 (2008).
- 41 S, S. *Effective Management of COPD in Primary Care: Challenges and Opportunities*. *AJMC* November (2018).
- 42 Valderas, J. M., Starfield, B., Sibbald, B., Salisburly, C. & Roland, M. Defining comorbidity: implications for understanding health and health services. *The Annals of Family Medicine* 7, 357-363 (2009).
- 43 Sethi, S. Infection as a comorbidity of COPD. *European Respiratory Journal* 35, 1209-1215 (2010).

- 44 Almirall, J. & Fortin, M. The coexistence of terms to describe the presence of multiple concurrent diseases. *Journal of comorbidity* 3, 4-9 (2013).
- 45 Drummond, M. B., Buist, A. S., Crapo, J. D., Wise, R. A. & Rennard, S. I. Chronic obstructive pulmonary disease: NHLBI workshop on the primary prevention of chronic lung diseases. *Annals of the American Thoracic Society* 11, S154-S160 (2014).
- 46 Sethi, S. & Murphy, T. F. Bacterial infection in chronic obstructive pulmonary disease in 2000: a state-of-the-art review. *Clin Microbiol Rev* 14, 336-363, doi:10.1128/cmr.14.2.336-363.2001 (2001).
- 47 Gibson, P. & Simpson, J. The overlap syndrome of asthma and COPD: what are its features and how important is it? *Thorax* 64, 728-735 (2009).
- 48 Zeki, A. A., Schivo, M., Chan, A., Albertson, T. E. & Louie, S. The asthma-COPD overlap syndrome: a common clinical problem in the elderly. *Journal of allergy* 2011 (2011).
- 49 Hunt, E. B., Sullivan, A., Galvin, J., MacSharry, J. & Murphy, D. M. Gastric aspiration and its role in airway inflammation. *The open respiratory medicine journal* 12, 1 (2018).
- 50 Chaouat, A., Naeije, R. & Weitzenblum, E. Pulmonary hypertension in COPD. *Eur Respir J* 32, 1371-1385, doi:10.1183/09031936.00015608 (2008).
- 51 Barnes, P. J. & Kharitonov, S. A. Exhaled nitric oxide: a new lung function test. *Thorax* 51, 233 (1996).
- 52 Hillas, G., Perlikos, F., Tsiligianni, I. & Tzanakis, N. Managing comorbidities in COPD. *International Journal of chronic obstructive pulmonary disease* 10, 95 (2015).
- 53 Stringer, W. & Marciniuk, D. The role of cardiopulmonary exercise testing (CPET) in pulmonary rehabilitation (PR) of chronic obstructive pulmonary disease (COPD) patients. *COPD: Journal of Chronic Obstructive Pulmonary Disease* 15, 621-631 (2018).
- 54 McKeough, Z. et al. Shuttle walk tests in people with COPD who demonstrate exercise-induced oxygen desaturation: An analysis of test repeatability and cardiorespiratory responses. *Chronic respiratory disease* 15, 131-137 (2018).
- 55 Chan, C., Yamabayashi, C., Syed, N., Kirkham, A. & Camp, P. G. Exercise telemonitoring and telerehabilitation compared with traditional cardiac and pulmonary rehabilitation: A systematic review and meta-analysis. *Physiotherapy Canada* 68, 242-251 (2016).
- 56 Bailey, K. L. The importance of the assessment of pulmonary function in COPD. *Medical Clinics* 96, 745-752 (2012).
- 57 Andreeva, E. et al. Spirometry is not enough to diagnose COPD in epidemiological studies: a follow-up study. *NPJ Prim Care Respir Med* 27, 62, doi:10.1038/s41533-017-0062-6 (2017).
- 58 Cerveri, I. et al. Underestimation of airflow obstruction among young adults using FEV<sub>1</sub>/FVC <70% as a fixed cut-off: a longitudinal evaluation of clinical and functional outcomes. *Thorax* 63, 1040-1045, doi:10.1136/thx.2008.095554 (2008).
- 59 Vollmer, W. M. et al. Comparison of spirometry criteria for the diagnosis of COPD: results from the BOLD study. *Eur Respir J* 34, 588-597, doi:10.1183/09031936.00164608 (2009).
- 60 Swanney, M. P. et al. Using the lower limit of normal for the FEV<sub>1</sub>/FVC ratio reduces the misclassification of airway obstruction. *Thorax* 63, 1046-1051, doi:10.1136/thx.2008.098483 (2008).
- 61 Guerra, S. Asthma and chronic obstructive pulmonary disease. *Curr Opin Allergy Clin Immunol* 9, 409-416, doi:10.1097/ACI.0b013e3283300baf (2009).
- 62 Bhatt, S. P. et al. Comparison of spirometric thresholds in diagnosing smoking-related airflow obstruction. *Thorax* 69, 409-414, doi:10.1136/thoraxjnl-2012-202810 (2014).
- 63 ANZ. The COPD-X Plan: Australian and New Zealand Guidelines for the management of Chronic Obstructive Pulmonary Disease <<https://copdx.org.au/>> (2020).

- 64 Chetta, A. & Olivieri, D. The COPD Assessment Test in the evaluation of chronic obstructive pulmonary disease exacerbations. *Expert Rev Respir Med* 6, 373-375, doi:10.1586/ers.12.37 (2012).
- 65 Munari, A. B. et al. Modified Medical Research Council Dyspnea Scale in GOLD Classification Better Reflects Physical Activities of Daily Living. *Respir Care* 63, 77-85, doi:10.4187/respcare.05636 (2018).
- 66 ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 166, 111-117, doi:10.1164/ajrccm.166.1.at1102 (2002).
- 67 AustralianLungFoundation. Incremental Shuttle Walk Test | Pulmonary Rehabilitation Toolkit, <<https://pulmonaryrehab.com.au/patient-assessment/assessing-exercise-capacity/incremental-shuttle-walking-test/>> (2020).
- 68 Agarwal, B., Shah, M., Andhare, N. & Mullerpatan, R. Incremental shuttle walk test: Reference values and predictive equation for healthy Indian adults. *Lung India* 33, 36-41, doi:10.4103/0970-2113.173056 (2016).
- 69 Singh, S. J., Morgan, M. D., Scott, S., Walters, D. & Hardman, A. E. Development of a shuttle walking test of disability in patients with chronic airways obstruction. *Thorax* 47, 1019-1024, doi:10.1136/thx.47.12.1019 (1992).
- 70 Singh, S. J., Morgan, M. D., Hardman, A. E., Rowe, C. & Bardsley, P. A. Comparison of oxygen uptake during a conventional treadmill test and the shuttle walking test in chronic airflow limitation. *Eur Respir J* 7, 2016-2020 (1994).
- 71 Miskoff, J. A., Dewan, A. & Chaudhri, M. Fractional Exhaled Nitric Oxide Testing: Diagnostic Utility in Asthma, Chronic Obstructive Pulmonary Disease, or Asthma-chronic Obstructive Pulmonary Disease Overlap Syndrome. *Cureus* 11, e4864, doi:10.7759/cureus.4864 (2019).
- 72 Dweik, R. A. et al. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FENO) for clinical applications. *Am J Respir Crit Care Med* 184, 602-615, doi:10.1164/rccm.9120-11ST (2011).
- 73 Tomikawa, M. et al. Asthma diagnosis and treatment - 1005. Optimization for the withdrawal of inhaled corticosteroid treatment by monitoring fractional exhaled nitric oxide (feno) and lung functions. *World Allergy Organ J* 6 Suppl 1, P5, doi:10.1186/1939-4551-6-s1-p5 (2013).
- 74 Asano, T. et al. Diagnostic utility of fractional exhaled nitric oxide in prolonged and chronic cough according to atopic status. *Allergol Int* 66, 344-350, doi:10.1016/j.alit.2016.08.015 (2017).
- 75 WHO. (2011).
- 76 Madan, A. Correlation between the levels of SpO<sub>2</sub> and PaO<sub>2</sub>. *Lung India : official organ of Indian Chest Society* 34, 307-308, doi:10.4103/lungindia.lungindia\_106\_17 (2017).
- 77 Sarkar, M., Niranjana, N. & Banyal, P. Mechanisms of hypoxemia. *Lung India* 34, 47-60, doi:10.4103/0970-2113.197116 (2017).
- 78 Burton Rose, T. P. *Clinical Physiology of Acid-Base and Electrolyte Disorders (Clinical Physiology of Acid Base & Electrolyte Disorders) 5th Edition* edn.
- 79 Fidkowski, C. & Helstrom, J. Diagnosing metabolic acidosis in the critically ill: bridging the anion gap, Stewart, and base excess methods. *Can J Anaesth* 56, 247-256, doi:10.1007/s12630-008-9037-y (2009).
- 80 David A. Kaufman. Interpretation of Arterial Blood Gases (ABGs) , <<https://www.thoracic.org/professionals/clinical-resources/critical-care/clinical-education/abgs.php>> (
- 81 Yalcin, A. D., Celik, B. & Yalcin, A. N. Omalizumab (anti-IgE) therapy in the asthma-COPD overlap syndrome (ACOS) and its effects on circulating cytokine levels. *Immunopharmacol Immunotoxicol* 38, 253-256, doi:10.3109/08923973.2016.1173057 (2016).

- 82 Lencu, C., Alexescu, T., Petrulea, M. & Lencu, M. Respiratory manifestations in endocrine diseases. *Clujul Med* 89, 459-463, doi:10.15386/cjmed-671 (2016).
- 83 Lane, R. & Adams, L. Metabolic acidosis and breathlessness during exercise and hypercapnia in man. *J Physiol* 461, 47-61, doi:10.1113/jphysiol.1993.sp019500 (1993).
- 84 Krapf, R., Beeler, I., Hertner, D. & Hulter, H. N. Chronic respiratory alkalosis. The effect of sustained hyperventilation on renal regulation of acid-base equilibrium. *N Engl J Med* 324, 1394-1401, doi:10.1056/nejm199105163242003 (1991).
- 85 Elif Torun Parmaksız, B. S., Gülşen Saraç, Nesrin Kiral, Ali Fidan, Sevda Cömert, Benan Çağlayan. Undiagnosed Hypothyroidism in Obstructive Sleep Apnea Syndrome *Journal of Turkish Sleep Medicine* 1, 11-14, doi:D0I: 10.4274/jtsm.02.003 (2015).
- 86 Fawzy, A. et al. Association of platelet count with all-cause mortality and risk of cardiovascular and respiratory morbidity in stable COPD. *Respir Res* 20, 86, doi:10.1186/s12931-019-1059-1 (2019).
- 87 Harrison, M. T. et al. Thrombocytosis is associated with increased short and long term mortality after exacerbation of chronic obstructive pulmonary disease: a role for antiplatelet therapy? *Thorax* 69, 609-615, doi:10.1136/thoraxjnl-2013-203996 (2014).
- 88 Blanco, I., Lara, B. & de Serres, F. Efficacy of alpha1-antitrypsin augmentation therapy in conditions other than pulmonary emphysema. *Orphanet J Rare Dis* 6, 14, doi:10.1186/1750-1172-6-14 (2011).
- 89 Al Ashry, H. S. & Strange, C. COPD in individuals with the PiMZ alpha-1 antitrypsin genotype. *Eur Respir Rev* 26, doi:10.1183/16000617.0068-2017 (2017).
- 90 Hylkema, M. N., Sterk, P. J., de Boer, W. I. & Postma, D. S. Tobacco use in relation to COPD and asthma. *Eur Respir J* 29, 438-445, doi:10.1183/09031936.00124506 (2007).
- 91 American Thoracic Society/European Respiratory Society statement: standards for the diagnosis and management of individuals with alpha-1 antitrypsin deficiency. *Am J Respir Crit Care Med* 168, 818-900, doi:10.1164/rccm.168.7.818 (2003).
- 92 Coxson, H. O. & Lam, S. Quantitative assessment of the airway wall using computed tomography and optical coherence tomography. *Proc Am Thorac Soc* 6, 439-443, doi:10.1513/pats.200904-015AW (2009).
- 93 Gupta, P. P., Yadav, R., Verma, M., Agarwal, D. & Kumar, M. Correlation between high-resolution computed tomography features and patients' characteristics in chronic obstructive pulmonary disease. *Ann Thorac Med* 3, 87-93, doi:10.4103/1817-1737.39676 (2008).
- 94 Young, K., Aspestrand, F. & Kolbenstvedt, A. High resolution CT and bronchography in the assessment of bronchiectasis. *Acta Radiol* 32, 439-441 (1991).
- 95 Jen, R., Li, Y., Owens, R. L. & Malhotra, A. Sleep in Chronic Obstructive Pulmonary Disease: Evidence Gaps and Challenges. *Can Respir J* 2016, 7947198, doi:10.1155/2016/7947198 (2016).
- 96 Pierce, A. K., Jarrett, C. E., Werkle, G., Jr. & Miller, W. F. Respiratory function during sleep in patients with chronic obstructive lung disease. *J Clin Invest* 45, 631-636, doi:10.1172/jci105378 (1966).
- 97 Cukic, V., Lovre, V., Dragisic, D. & Ustamujic, A. Asthma and Chronic Obstructive Pulmonary Disease (COPD) - Differences and Similarities. *Mater Sociomed* 24, 100-105, doi:10.5455/msm.2012.24.100-105 (2012).
- 98 Gao, J. & Wu, F. Association between fractional exhaled nitric oxide, sputum induction and peripheral blood eosinophil in uncontrolled asthma. *Allergy Asthma Clin Immunol* 14, 21, doi:10.1186/s13223-018-0248-7 (2018).
- 99 Platts-Mills, T. A. E., Schuyler, A. J., Erwin, E. A., Commins, S. P. & Woodfolk, J. A. IgE in the diagnosis and treatment of allergic disease. *J Allergy Clin Immunol* 137, 1662-

- 1670, doi:10.1016/j.jaci.2016.04.010 (2016).
- 100 Gao, J. et al. Correlation between fractional exhaled nitric oxide and sputum eosinophilia in exacerbations of COPD. *Int J Chron Obstruct Pulmon Dis* 12, 1287-1293, doi:10.2147/copd.S134998 (2017).
- 101 Donohue, J. F. & Jain, N. Exhaled nitric oxide to predict corticosteroid responsiveness and reduce asthma exacerbation rates. *Respir Med* 107, 943-952, doi:10.1016/j.rmed.2013.02.018 (2013).
- 102 Beg, M. F., Alzogaibi, M. A., Abba, A. A. & Habib, S. S. Exhaled nitric oxide in stable chronic obstructive pulmonary disease. *Ann Thorac Med* 4, 65-70, doi:10.4103/1817-1737.44649 (2009).
- 103 Deng, D. D., Zhou, A. Y., Shuang, Q. C. & Chen, P. [The value of fractionated exhaled nitric oxide in the diagnosis of asthma-chronic obstructive pulmonary disease overlap syndrome]. *Zhonghua Jie He He Hu Xi Za Zhi* 40, 98-101, doi:10.3760/cma.j.isn.1001-0939.2017.02.004 (2017).
- 104 Fabbri, L. M., Beghé, B. & Papi, A. Blood eosinophils for the management of COPD patients? *Lancet Respir Med* 6, 807-808, doi:10.1016/s2213-2600(18)30417-x (2018).
- 105 Rodrigo, G. J. et al. LABA/LAMA combinations versus LAMA monotherapy or LABA/ICS in COPD: a systematic review and meta-analysis. *Int J Chron Obstruct Pulmon Dis* 12, 907-922, doi:10.2147/copd.S130482 (2017).
- 106 Bandyopadhyay, A. et al. Usefulness of induced sputum eosinophil count to assess severity and treatment outcome in asthma patients. *Lung India* 30, 117-123, doi:10.4103/0970-2113.110419 (2013).
- 107 Oliver, B. et al. COPD treatment choices based on blood eosinophils: are we there yet? *Breathe (Sheff)* 15, 318-323, doi:10.1183/20734735.0254-2019 (2019).
- 108 Kumar, R. M., Pajanivel, R., Koteeswaran, G., Menon, S. K. & Charles, P. M. Correlation of total serum immunoglobulin E level, sputum, and peripheral eosinophil count in assessing the clinical severity in bronchial asthma. *Lung India* 34, 256-261, doi:10.4103/lungindia.lungindia\_73\_16 (2017).
- 109 Clinic, M. Fractional exhaled nitric oxide moderately accurate to diagnose asthma., <[https://medicalxpress.com/news/2018-01-fractional-exhaled-nitric-oxide-moderately.html?utm\\_source=TrendMD&utm\\_medium=cpc&utm\\_campaign=MedicalXpress](https://medicalxpress.com/news/2018-01-fractional-exhaled-nitric-oxide-moderately.html?utm_source=TrendMD&utm_medium=cpc&utm_campaign=MedicalXpress)> (2020).
- 110 Rogliani, P., Ora, J., Puxeddu, E. & Cazzola, M. Airflow obstruction: is it asthma or is it COPD? *Int J Chron Obstruct Pulmon Dis* 11, 3007-3013, doi:10.2147/copd.S54927 (2016).
- 111 Miller, W. T., Jr., Chatzkel, J. & Hewitt, M. G. Expiratory air trapping on thoracic computed tomography. A diagnostic subclassification. *Ann Am Thorac Soc* 11, 874-881, doi:10.1513/AnnalsATS.201311-3900C (2014).
- 112 Parveen T , B. N., Begum S Allergen Skin Test Reactivity and Serum Total IgE Level in Adult Bronchial Asthmatic Patients. *J Bangladesh Soc Physiol.* 4, 1-6 (2009).
- 113 Martinez-Garcia, M. A. & Miravittles, M. Bronchiectasis in COPD patients: more than a comorbidity? *Int J Chron Obstruct Pulmon Dis* 12, 1401-1411, doi:10.2147/copd.S132961 (2017).
- 114 McDonnell, M. J. et al. Comorbidities and the risk of mortality in patients with bronchiectasis: an international multicentre cohort study. *Lancet Respir Med* 4, 969-979, doi:10.1016/s2213-2600(16)30320-4 (2016).
- 115 Chen, Y. H. & Sun, Y. C. Bronchiectasis as a Comorbidity of Chronic Obstructive Pulmonary Disease: Implications and Future Research. *Chin Med J (Engl)* 129, 2017-2019, doi:10.4103/0366-6999.189071 (2016).
- 116 Jin, J. et al. Factors associated with bronchiectasis in patients with moderate-severe chronic obstructive pulmonary disease. *Medicine (Baltimore)* 95, e4219, doi:10.1097/md.0000000000004219 (2016).

- 117 Jin, J., Li, S., Yu, W., Liu, X. & Sun, Y. Emphysema and bronchiectasis in COPD patients with previous pulmonary tuberculosis: computed tomography features and clinical implications. *Int J Chron Obstruct Pulmon Dis* 13, 375-384, doi:10.2147/copd.S152447 (2018).
- 118 Singh, D., Edwards, L., Tal-Singer, R. & Rennard, S. Sputum neutrophils as a biomarker in COPD: findings from the ECLIPSE study. *Respir Res* 11, 77, doi:10.1186/1465-9921-11-77 (2010).
- 119 Imaging of Bronchiectasis: Bronchiectasis Toolbox, <<https://bronchiectasis.com.au/bronchiectasis/diagnosis-2/radiology>> (2020).
- 120 Guan, W. J., Chen, R. C. & Zhong, N. S. The bronchiectasis severity index and FACED score for bronchiectasis. *Eur Respir J* 47, 382-384, doi:10.1183/13993003.01717-2015 (2016).
- 121 Fujimoto, K. et al. Airway inflammation during stable and acutely exacerbated chronic obstructive pulmonary disease. *Eur Respir J* 25, 640-646, doi:10.1183/09031936.05.00047504 (2005).
- 122 Fuschillo, S., Martucci, M., Donner, C. F. & Balzano, G. Airway bacterial colonization: the missing link between COPD and cardiovascular events? *Respir Med* 106, 915-923, doi:10.1016/j.rmed.2012.03.023 (2012).
- 123 Zhang, M. et al. Relevance of lower airway bacterial colonization, airway inflammation, and pulmonary function in the stable stage of chronic obstructive pulmonary disease. *Eur J Clin Microbiol Infect Dis* 29, 1487-1493, doi:10.1007/s10096-010-1027-7 (2010).
- 124 Patel, I. S. et al. Bronchiectasis, exacerbation indices, and inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 170, 400-407, doi:10.1164/rccm.200305-6480C (2004).
- 125 Gevenois, P. A., de Maertelaer, V., De Vuyst, P., Zanen, J. & Yernault, J. C. Comparison of computed density and macroscopic morphometry in pulmonary emphysema. *Am J Respir Crit Care Med* 152, 653-657, doi:10.1164/ajrccm.152.2.7633722 (1995).
- 126 Gevenois, P. A. et al. Comparison of computed density and microscopic morphometry in pulmonary emphysema. *Am J Respir Crit Care Med* 154, 187-192, doi:10.1164/ajrccm.154.1.8680679 (1996).
- 127 Coban, H. & Gungen, A. C. Is There a Correlation between New Scoring Systems and Systemic Inflammation in Stable Bronchiectasis? *Can Respir J* 2017, 9874068, doi:10.1155/2017/9874068 (2017).
- 128 Habesoglu, M. A., Tercan, F., Ozkan, U. & Fusun, E. O. Effect of radiological extent and severity of bronchiectasis on pulmonary function. *Multidiscip Respir Med* 6, 284-290, doi:10.1186/2049-6958-6-5-284 (2011).
- 129 Khair, O. A. et al. Effect of Haemophilus influenzae endotoxin on the synthesis of IL-6, IL-8, TNF-alpha and expression of ICAM-1 in cultured human bronchial epithelial cells. *Eur Respir J* 7, 2109-2116, doi:10.1183/09031936.94.07122109 (1994).
- 130 MacNee, W. Pathology, pathogenesis, and pathophysiology. *BMJ : British Medical Journal* 332, 1202-1204 (2006).
- 131 Sverzellati, N. et al. New insights on COPD imaging via CT and MRI. *Int J Chron Obstruct Pulmon Dis* 2, 301-312 (2007).
- 132 Dou, S. et al. High prevalence of bronchiectasis in emphysema-predominant COPD patients. *Int J Chron Obstruct Pulmon Dis* 13, 2041-2047, doi:10.2147/copd.S163243 (2018).
- 133 Badawy MS, E. Q. A., Mohamadeen HA. Alpha1 antitrypsin deficiency in non cystic fibrosis bronchiectasis. *Egyptian Journal of Chest Diseases and Tuberculosis* 2, 62 (2013).
- 134 Pomares, X. et al. Long-term azithromycin therapy in patients with severe COPD and repeated exacerbations. *Int J Chron Obstruct Pulmon Dis* 6, 449-456, doi:10.2147/

- copd.S23655 (2011).
- 135 Dal Negro, R., Micheletto, C., Tognella, S., Visconti, M. & Turati, C. Tobramycin Nebulizer Solution in severe COPD patients colonized with *Pseudomonas aeruginosa*: effects on bronchial inflammation. *Adv Ther* 25, 1019-1030, doi:10.1007/s12325-008-0105-2 (2008).
- 136 Chow, E. J., Doyle, J. D. & Uyeki, T. M. Influenza virus-related critical illness: prevention, diagnosis, treatment. *Crit Care* 23, 214, doi:10.1186/s13054-019-2491-9 (2019).
- 137 Froes, F., Roche, N. & Blasi, F. Pneumococcal vaccination and chronic respiratory diseases. *Int J Chron Obstruct Pulmon Dis* 12, 3457-3468, doi:10.2147/copd.S140378 (2017).
- 138 Braido, F., Tarantini, F., Ghiglione, V., Melioli, G. & Canonica, G. W. Bacterial lysate in the prevention of acute exacerbation of COPD and in respiratory recurrent infections. *Int J Chron Obstruct Pulmon Dis* 2, 335-345 (2007).
- 139 Casanova, C. et al. Increased gastro-oesophageal reflux disease in patients with severe COPD. *Eur Respir J* 23, 841-845, doi:10.1183/09031936.04.00107004 (2004).
- 140 Lee, A. L. & Goldstein, R. S. Gastroesophageal reflux disease in COPD: links and risks. *Int J Chron Obstruct Pulmon Dis* 10, 1935-1949, doi:10.2147/copd.S77562 (2015).
- 141 Shin, J. M. et al. Molecular mechanisms in therapy of acid-related diseases. *Cell Mol Life Sci* 65, 264-281, doi:10.1007/s00018-007-7249-x (2008).
- 142 Huang, C., Liu, Y. & Shi, G. A systematic review with meta-analysis of gastroesophageal reflux disease and exacerbations of chronic obstructive pulmonary disease. *BMC Pulm Med* 20, 2, doi:10.1186/s12890-019-1027-z (2020).
- 143 Broers, C., Tack, J. & Pauwels, A. Review article: gastro-oesophageal reflux disease in asthma and chronic obstructive pulmonary disease. *Aliment Pharmacol Ther* 47, 176-191, doi:10.1111/apt.14416 (2018).
- 144 Herzig, S. J., Howell, M. D., Ngo, L. H. & Marcantonio, E. R. Acid-suppressive medication use and the risk for hospital-acquired pneumonia. *Jama* 301, 2120-2128, doi:10.1001/jama.2009.722 (2009).
- 145 Lin, W. L., Muo, C. S., Lin, W. C., Hsieh, Y. W. & Kao, C. H. Association of Increased Risk of Pneumonia and Using Proton Pump Inhibitors in Patients With Type II Diabetes Mellitus. *Dose Response* 17, 1559325819843383, doi:10.1177/1559325819843383 (2019).
- 146 AH., M. Airway reflux as a cause of respiratory disease. *Breathe* 9, 256-266 (2013).
- 147 Gaude, G. S. Pulmonary manifestations of gastroesophageal reflux disease. *Annals of Thoracic Medicine* 4, 115 (2009).
- 148 Mokhlesi, B. et al. Increased prevalence of gastroesophageal reflux symptoms in patients with COPD. *Chest* 119, 1043-1048, doi:10.1378/chest.119.4.1043 (2001).
- 149 Kiljander, T. O. et al. Effects of esomeprazole 40 mg twice daily on asthma: a randomized placebo-controlled trial. *Am J Respir Crit Care Med* 173, 1091-1097, doi:10.1164/rccm.200507-11670C (2006).
- 150 Kikuchi S, N. Y., Tajiri T, Watanabe N. (2018).
- 151 Hochhegger, B. et al. Imaging in idiopathic pulmonary fibrosis: diagnosis and mimics. *Clinics (Sao Paulo)* 74, e225, doi:10.6061/clinics/2019/e225 (2019).
- 152 Oikonomou, A. et al. Pulmonary fibrosis and emphysema: Is the emphysema type associated with the pattern of fibrosis? *World J Radiol* 7, 294-305, doi:10.4329/wjr.v7.i9.294 (2015).
- 153 Hashemi, S. J., Javad Moosavi, S. A., Idani, E., Javaherforoushzadeh, A. & Raji, H. Association of esophageal dilatation in chest CT scan with gastroesophageal reflux disease: A case control study. *Monaldi Arch Chest Dis* 88, 914, doi:10.4081/monaldi.2018.914 (2018).
- 154 Neto, R. M. L., Herbella, F. A. M., Schlottmann, F. & Patti, M. G. Does DeMeester score

- still define GERD? *Dis Esophagus* 32, doi:10.1093/dote/doy118 (2019).
- 155 Owens, R. L. & Malhotra, A. Sleep-disordered breathing and COPD: the overlap syndrome. *Respir Care* 55, 1333-1344; discussion 1344-1336 (2010).
- 156 Soler, X. et al. High Prevalence of Obstructive Sleep Apnea in Patients with Moderate to Severe Chronic Obstructive Pulmonary Disease. *Ann Am Thorac Soc* 12, 1219-1225, doi:10.1513/AnnalsATS.201407-3360C (2015).
- 157 Cormick, W., Olson, L. G., Hensley, M. J. & Saunders, N. A. Nocturnal hypoxaemia and quality of sleep in patients with chronic obstructive lung disease. *Thorax* 41, 846-854, doi:10.1136/thx.41.11.846 (1986).
- 158 Klink, M. E., Dodge, R. & Quan, S. F. The relation of sleep complaints to respiratory symptoms in a general population. *Chest* 105, 151-154, doi:10.1378/chest.105.1.151 (1994).
- 159 Ezzie, M. E., Parsons, J. P. & Mastronarde, J. G. Sleep and Obstructive Lung Diseases. *Sleep Med Clin* 3, 505-515, doi:10.1016/j.jsmc.2008.07.003 (2008).
- 160 Douglas, N. J. et al. Hypoxic ventilatory response decreases during sleep in normal men. *Am Rev Respir Dis* 125, 286-289, doi:10.1164/arrd.1982.125.3.286 (1982).
- 161 Douglas, N. J., White, D. P., Weil, J. V., Pickett, C. K. & Zwillich, C. W. Hypercapnic ventilatory response in sleeping adults. *Am Rev Respir Dis* 126, 758-762, doi:10.1164/arrd.1982.126.5.758 (1982).
- 162 Hudgel, D. W., Martin, R. J., Johnson, B. & Hill, P. Mechanics of the respiratory system and breathing pattern during sleep in normal humans. *J Appl Physiol Respir Environ Exerc Physiol* 56, 133-137, doi:10.1152/jappl.1984.56.1.133 (1984).
- 163 Fletcher, E. C., Schaaf, J. W., Miller, J. & Fletcher, J. G. Long-term cardiopulmonary sequelae in patients with sleep apnea and chronic lung disease. *Am Rev Respir Dis* 135, 525-533, doi:10.1164/arrd.1987.135.3.525 (1987).
- 164 Lewis, C. A., Fergusson, W., Eaton, T., Zeng, I. & Kolbe, J. Isolated nocturnal desaturation in COPD: prevalence and impact on quality of life and sleep. *Thorax* 64, 133-138, doi:10.1136/thx.2007.088930 (2009).
- 165 Mulloy, E., Fitzpatrick, M., Bourke, S., O'Regan, A. & McNicholas, W. T. Oxygen desaturation during sleep and exercise in patients with severe chronic obstructive pulmonary disease. *Respir Med* 89, 193-198, doi:10.1016/0954-6111(95)90247-3 (1995).
- 166 Mulloy, E. & McNicholas, W. T. Ventilation and gas exchange during sleep and exercise in severe COPD. *Chest* 109, 387-394, doi:10.1378/chest.109.2.387 (1996).
- 167 Fletcher, E. C. et al. Survival in COPD patients with a daytime PaO<sub>2</sub> greater than 60 mm Hg with and without nocturnal oxyhemoglobin desaturation. *Chest* 101, 649-655, doi:10.1378/chest.101.3.649 (1992).
- 168 Celli, B. R. & MacNee, W. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J* 23, 932-946, doi:10.1183/09031936.04.00014304 (2004).
- 169 Spicuzza, L., Caruso, D. & Di Maria, G. Obstructive sleep apnoea syndrome and its management. *Ther Adv Chronic Dis* 6, 273-285, doi:10.1177/2040622315590318 (2015).
- 170 Kessler, R. et al. Pulmonary hypertension in the obstructive sleep apnoea syndrome: prevalence, causes and therapeutic consequences. *Eur Respir J* 9, 787-794, doi:10.1183/09031936.96.09040787 (1996).
- 171 Resta, O. et al. Prescription of nCPAP and nBIPAP in obstructive sleep apnoea syndrome: Italian experience in 105 subjects. A prospective two centre study. *Respir Med* 92, 820-827, doi:10.1016/s0954-6111(98)90383-3 (1998).
- 172 Roy TM, M. Z. *Respiratory Acidosis: Practice Essentials*, <emedicine.medscape.com> (2019).
- 173 Cukic, V. The changes of arterial blood gases in COPD during four-year period. *Med Arch* 68, 14-18, doi:10.5455/medarh.2014.68.14-18 (2014).



- 174 Sun, W. L. et al. Impact of obstructive sleep apnea on pulmonary hypertension in patients with chronic obstructive pulmonary disease. *Chin Med J (Engl)* 132, 1272-1282, doi:10.1097/cm9.000000000000247 (2019).
- 175 Cade, B. E. et al. Associations of variants in the hexokinase 1 and interleukin 18 receptor regions with oxyhemoglobin saturation during sleep. *PLoS Genet* 15, e1007739, doi:10.1371/journal.pgen.1007739 (2019).
- 176 Ascha, M., Renapurkar, R. D. & Tonelli, A. R. A review of imaging modalities in pulmonary hypertension. *Ann Thorac Med* 12, 61-73, doi:10.4103/1817-1737.203742 (2017).
- 177 Lynch, D. A. et al. CT-Definable Subtypes of Chronic Obstructive Pulmonary Disease: A Statement of the Fleischner Society. *Radiology* 277, 192-205, doi:10.1148/radiol.2015141579 (2015).
- 178 D'Alto, M. et al. Accuracy and precision of echocardiography versus right heart catheterization for the assessment of pulmonary hypertension. *Int J Cardiol* 168, 4058-4062, doi:10.1016/j.ijcard.2013.07.005 (2013).
- 179 Greiner, S. et al. Reliability of noninvasive assessment of systolic pulmonary artery pressure by Doppler echocardiography compared to right heart catheterization: analysis in a large patient population. *J Am Heart Assoc* 3, doi:10.1161/jaha.114.001103 (2014).
- 180 Hammerstingl, C. et al. Diagnostic value of echocardiography in the diagnosis of pulmonary hypertension. *PLoS One* 7, e38519, doi:10.1371/journal.pone.0038519 (2012).
- 181 Rosenthal, L. D. & Dolan, D. C. The Epworth sleepiness scale in the identification of obstructive sleep apnea. *J Nerv Ment Dis* 196, 429-431, doi:10.1097/NMD.0b013e31816ff3bf (2008).
- 182 Chung, F., Abdullah, H. R. & Liao, P. STOP-Bang Questionnaire: A Practical Approach to Screen for Obstructive Sleep Apnea. *Chest* 149, 631-638, doi:10.1378/chest.15-0903 (2016).
- 183 Chung, F. et al. High STOP-Bang score indicates a high probability of obstructive sleep apnoea. *Br J Anaesth* 108, 768-775, doi:10.1093/bja/aes022 (2012).
- 184 Ko, F. W. & Hui, D. S. Outdoor air pollution: impact on chronic obstructive pulmonary disease patients. *Curr Opin Pulm Med* 15, 150-157, doi:10.1097/MCP.0b013e32832185ee (2009).
- 185 Liu, Y., Lee, K., Perez-Padilla, R., Hudson, N. L. & Mannino, D. M. Outdoor and indoor air pollution and COPD-related diseases in high- and low-income countries. *Int J Tuberc Lung Dis* 12, 115-127 (2008).
- 186 Health effects of outdoor air pollution. Part 2. Committee of the Environmental and Occupational Health Assembly of the American Thoracic Society. *Am J Respir Crit Care Med* 153, 477-498, doi:10.1164/ajrccm.153.2.8564086 (1996).
- 187 Jiang, X. Q., Mei, X. D. & Feng, D. Air pollution and chronic airway diseases: what should people know and do? *J Thorac Dis* 8, E31-40, doi:10.3978/j.issn.2072-1439.2015.11.50 (2016).
- 188 Schikowski, T. et al. Ambient air pollution: a cause of COPD? *Eur Respir J* 43, 250-263, doi:10.1183/09031936.00100112 (2014).
- 189 Li, J. et al. Major air pollutants and risk of COPD exacerbations: a systematic review and meta-analysis. *Int J Chron Obstruct Pulmon Dis* 11, 3079-3091, doi:10.2147/copd.S122282 (2016).
- 190 Hansel, N. N., McCormack, M. C. & Kim, V. The Effects of Air Pollution and Temperature on COPD. *COPD* 13, 372-379, doi:10.3109/15412555.2015.1089846 (2016).
- 191 McCormack, M. C. et al. Respiratory Effects of Indoor Heat and the Interaction with Air Pollution in Chronic Obstructive Pulmonary Disease. *Ann Am Thorac Soc* 13, 2125-2131, doi:10.1513/AnnalsATS.201605-3290C (2016).

- 192 Michelozzi, P. et al. High temperature and hospitalizations for cardiovascular and  
respiratory causes in 12 European cities. *Am J Respir Crit Care Med* 179, 383-389,  
doi:10.1164/rccm.200802-2170C (2009).
- 193 Lin, S. et al. Extreme high temperatures and hospital admissions for respiratory and  
cardiovascular diseases. *Epidemiology*, 738-746 (2009).
- 194 Tseng, C. M. et al. The effect of cold temperature on increased exacerbation of  
chronic obstructive pulmonary disease: a nationwide study. *PLoS One* 8, e57066,  
doi:10.1371/journal.pone.0057066 (2013).
- 195 D'Amato, M. et al. The impact of cold on the respiratory tract and its consequenc-  
es to respiratory health. *Clin Transl Allergy* 8, 20, doi:10.1186/s13601-018-0208-9  
(2018).
- 196 Shen, Y. et al. Management of airway mucus hypersecretion in chronic airway in-  
flammatory disease: Chinese expert consensus (English edition). *Int J Chron Obstruct  
Pulmon Dis* 13, 399-407, doi:10.2147/copd.S144312 (2018).
- 197 Cavallès, A. et al. Comorbidities of COPD. *Eur Respir Rev* 22, 454-475,  
doi:10.1183/09059180.00008612 (2013).
- 198 Panetta, N. L., Krachman, S. & Chatila, W. M. Chronic obstructive pulmonary disease  
and its comorbidities. *Panminerva Med* 51, 115-123 (2009).
- 199 Aryal, S., Diaz-Guzman, E. & Mannino, D. M. Influence of sex on chronic obstructive  
pulmonary disease risk and treatment outcomes. *Int J Chron Obstruct Pulmon Dis* 9,  
1145-1154, doi:10.2147/copd.S54476 (2014).
- 200 Morgan, A. D., Rothnie, K. J., Bhaskaran, K., Smeeth, L. & Quint, J. K. Chronic ob-  
structive pulmonary disease and the risk of 12 cardiovascular diseases: a popu-  
lation-based study using UK primary care data. *Thorax* 73, 877-879, doi:10.1136/  
thoraxjnl-2017-210865 (2018).
- 201 Morgan, A. D., Zakeri, R. & Quint, J. K. Defining the relationship between COPD  
and CVD: what are the implications for clinical practice? *Ther Adv Respir Dis* 12,  
1753465817750524, doi:10.1177/1753465817750524 (2018).
- 202 Macnee, W., Maclay, J. & McAllister, D. Cardiovascular injury and repair in chronic  
obstructive pulmonary disease. *Proceedings of the American Thoracic Society* 5,  
824-833, doi:10.1513/pats.200807-071TH (2008).
- 203 Vivodtzev, I. et al. Arterial stiffness in COPD. *Chest* 145, 861-875, doi:10.1378/  
chest.13-1809 (2014).
- 204 Clinic, M. Heart arrhythmia - Diagnosis and treatment <[www.mayoclinic.org](http://www.mayoclinic.org)> diag-  
no-sis-treatment > drc-20350674> (
- 205 Montuschi, P. Pharmacological treatment of chronic obstructive pulmonary dis-  
ease. *International journal of chronic obstructive pulmonary disease* 1, 409-423,  
doi:10.2147/copd.2006.1.4.409 (2006).
- 206 Albert, R. K. & Schuller, J. L. Macrolide antibiotics and the risk of cardiac arrhythmias.  
*Am J Respir Crit Care Med* 189, 1173-1180, doi:10.1164/rccm.201402-0385CI  
(2014).
- 207 Almalki, Z. S. & Guo, J. J. Cardiovascular events and safety outcomes associated with  
azithromycin therapy: a meta-analysis of randomized controlled trials. *Am Health  
Drug Benefits* 7, 318-328 (2014).
- 208 Hannink, J. D., van Helvoort, H. A., Dekhuijzen, P. N. & Heijdra, Y. F. Heart failure  
and COPD: partners in crime? *Respirology* 15, 895-901, doi:10.1111/j.1440-  
1843.2010.01776.x (2010).
- 209 Mahendra, M., S, S. K., Desai, N., Bs, J. & Pa, M. Evaluation for airway obstruction  
in adult patients with stable ischemic heart disease. *Indian Heart J* 70, 266-271,  
doi:10.1016/j.ihj.2017.08.003 (2018).
- 210 Hawkins, N. M., Khosla, A., Virani, S. A., McMurray, J. J. & FitzGerald, J. M. B-type  
natriuretic peptides in chronic obstructive pulmonary disease: a systematic review.

- BMC Pulm Med 17, 11, doi:10.1186/s12890-016-0345-7 (2017).
- 211 Weber, M. & Hamm, C. Role of B-type natriuretic peptide (BNP) and NT-proBNP in  
clinical routine. *Heart* 92, 843-849, doi:10.1136/hrt.2005.071233 (2006).
- 212 McCullough, P.A. et al. B-type natriuretic peptide and clinical judgment in emergency  
diagnosis of heart failure: analysis from Breathing Not Properly (BNP) Multinational  
Study. *Circulation* 106, 416-422, doi:10.1161/01.cir.0000025242.79963.4c (2002).
- 213 FA, M. Empiric NT-proBNP Cut-Points for Diagnosing Acute Heart Failure. *NEJM*  
*JOURNAL WATCH* (2018).
- 214 ACC. Left Ventricular Ejection Fraction LVEF Assessment., <www. acc.org › clinical-  
toolkits › heart-failure-practice-solutions> (2017).
- 215 de Lucas-Ramos, P. et al. Chronic obstructive pulmonary disease as a cardiovascular  
risk factor. Results of a case-control study (CONSISTE study). *Int J Chron Obstruct*  
*Pulmon Dis* 7, 679-686, doi:10.2147/copd.S36222 (2012).
- 216 Etmnan, M., Jafari, S., Carleton, B. & FitzGerald, J. M. Beta-blocker use and  
COPD mortality: a systematic review and meta-analysis. *BMC Pulm Med* 12, 48,  
doi:10.1186/1471-2466-12-48 (2012).
- 217 Dransfield, M. T. et al. -Blocker Therapy and Clinical Outcomes in Patients with  
Moderate Chronic Obstructive Pulmonary Disease and Heightened Cardiovascular  
Risk. An Observational Substudy of SUMMIT. *Ann Am Thorac Soc* 15, 608-614,  
doi:10.1513/AnnalsATS.201708-6260C (2018).
- 218 Sarkar, M., Bhardwaj, R., Madabhavi, I. & Khatana, J. Osteoporosis in chronic ob-  
structive pulmonary disease. *Clin Med Insights Circ Respir Pulm Med* 9, 5-21,  
doi:10.4137/CCRPM.S22803 (2015).
- 219 Graat-Verboom, L. et al. Current status of research on osteoporosis in COPD: a sys-  
tematic review. *Eur Respir J* 34, 209-218, doi:10.1183/09031936.50130408 (2009).
- 220 Pascual-Guardia, S. et al. Hospital admissions and mortality in patients with COPD  
exacerbations and vertebral body compression fractures. *Int J Chron Obstruct Pulm-*  
*on Dis* 12, 1837-1845, doi:10.2147/copd.S129213 (2017).
- 221 MacLean, C. et al. Systematic review: comparative effectiveness of treatments to  
prevent fractures in men and women with low bone density or osteoporosis. *Ann In-*  
*tern Med* 148, 197-213, doi:10.7326/0003-4819-148-3-200802050-00198 (2008).
- 222 Islam AKMMA, H. M., Rahman KMc, Asaduzzaman Md, Rahim MAe, Zaman Sf, Islam  
MRg, Jesmin Hh, Yeasmin Li, . Vitamin D status in Bangladeshi subjects: a laboratory  
based study. *BIRDEM Med J* 9, 202-206 (2019).
- 223 Chen, X., Mao, G. & Leng, S. X. Frailty syndrome: an overview. *Clin Interv Aging* 9,  
433-441, doi:10.2147/cia.S45300 (2014).
- 224 Fried, L. P. et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci*  
*Med Sci* 56, M146-156, doi:10.1093/gerona/56.3.m146 (2001).
- 225 Morley, J. E., Malmstrom, T. K. & Miller, D. K. A simple frailty questionnaire (FRAIL)  
predicts outcomes in middle aged African Americans. *J Nutr Health Aging* 16, 601-  
608, doi:10.1007/s12603-012-0084-2 (2012).
- 226 Cesari, M. et al. A self-reported screening tool for detecting community-dwelling  
older persons with frailty syndrome in the absence of mobility disability: the FiND  
questionnaire. *PLoS one* 9, e101745 (2014).
- 227 Gale, N. S. et al. Frailty: A global measure of the multisystem impact of COPD. *Chron*  
*Respir Dis* 15, 347-355, doi:10.1177/1479972317752763 (2018).
- 228 Maddocks, M. et al. Physical frailty and pulmonary rehabilitation in COPD: a prospec-  
tive cohort study. *Thorax* 71, 988-995, doi:10.1136/thoraxjnl-2016-208460 (2016).
- 229 Guan, C. & Niu, H. Frailty assessment in older adults with chronic obstructive respi-  
ratory diseases. *Clin Interv Aging* 13, 1513-1524, doi:10.2147/cia.S173239 (2018).
- 230 Zhou, J. et al. Association of vitamin D deficiency and frailty: A systematic review and  
meta-analysis. *Maturitas* 94, 70-76, doi:10.1016/j.maturitas.2016.09.003 (2016).

- 231 Michelon, E. et al. Vitamin and carotenoid status in older women: associations with the frailty syndrome. *J Gerontol A Biol Sci Med Sci* 61, 600-607, doi:10.1093/gerona/61.6.600 (2006).
- 232 Keranis, E. et al. Impact of dietary shift to higher-antioxidant foods in COPD: a randomised trial. *Eur Respir J* 36, 774-780, doi:10.1183/09031936.00113809 (2010).
- 233 Peh, H. Y. et al. Vitamin E isoform  $\gamma$ -tocotrienol protects against emphysema in cigarette smoke-induced COPD. *Free Radic Biol Med* 110, 332-344, doi:10.1016/j.freeradbiomed.2017.06.023 (2017).
- 234 Matsuyama, W. et al. Effects of omega-3 polyunsaturated fatty acids on inflammatory markers in COPD. *Chest* 128, 3817-3827, doi:10.1378/chest.128.6.3817 (2005).
- 235 Sugawara, K. et al. Effects of nutritional supplementation combined with low-intensity exercise in malnourished patients with COPD. *Respir Med* 104, 1883-1889, doi:10.1016/j.rmed.2010.05.008 (2010).
- 236 Fuld, J. P. et al. Creatine supplementation during pulmonary rehabilitation in chronic obstructive pulmonary disease. *Thorax* 60, 531-537, doi:10.1136/thx.2004.030452 (2005).
- 237 Page, S. T. et al. Exogenous testosterone (T) alone or with finasteride increases physical performance, grip strength, and lean body mass in older men with low serum T. *J Clin Endocrinol Metab* 90, 1502-1510, doi:10.1210/jc.2004-1933 (2005).
- 238 Sharma, S. et al. Anabolic steroids in COPD: a review and preliminary results of a randomized trial. *Chron Respir Dis* 5, 169-176, doi:10.1177/1479972308092350 (2008).
- 239 Lo, Y. L. et al. Dietary Pattern Associated with Frailty: Results from Nutrition and Health Survey in Taiwan. *J Am Geriatr Soc* 65, 2009-2015, doi:10.1111/jgs.14972 (2017).
- 240 Oktaviana, J., Zanker, J., Vogrin, S. & Duque, G. The Effect of  $\beta$ -hydroxy- $\beta$ -methylbutyrate (HMB) on Sarcopenia and Functional Frailty in Older Persons: A Systematic Review. *J Nutr Health Aging* 23, 145-150, doi:10.1007/s12603-018-1153-y (2019).
- 241 Crocker, T. et al. Physical rehabilitation for older people in long-term care. *Cochrane Database Syst Rev*, Cd004294, doi:10.1002/14651858.CD004294.pub3 (2013).
- 242 Corhay, J.-L., Dang, D. N., Van Cauwenberge, H. & Louis, R. Pulmonary rehabilitation and COPD: providing patients a good environment for optimizing therapy. *International journal of chronic obstructive pulmonary disease* 9, 27 (2014).
- 243 Goërtz, Y. M. J. et al. Fatigue is highly prevalent in patients with COPD and correlates poorly with the degree of airflow limitation. *Ther Adv Respir Dis* 13, 1753466619878128, doi:10.1177/1753466619878128 (2019).
- 244 Medina-Mirapeix, F. et al. Patterns and predictors of exhaustion episodes in patients with stable COPD: A longitudinal study. *Int J Clin Pract* 72, e13068, doi:10.1111/ijcp.13068 (2018).
- 245 Deshmukh, A. & Schuller, D. Cough syncope in a patient with severe chronic obstructive pulmonary disease. *Respiratory Medicine CME* 1, 120-122 (2008).
- 246 Hakamy, A., Bolton, C. E., Gibson, J. E. & McKeever, T. M. Risk of fall in patients with COPD. *Thorax* 73, 1079-1080, doi:10.1136/thoraxjnl-2017-211008 (2018).
- 247 Crisan, A. F., Oancea, C., Timar, B., Fira-Mladinescu, O. & Tudorache, V. (Eur Respiratory Soc, 2015).
- 248 Beauchamp, M. K. et al. Pulmonary Rehabilitation With Balance Training for Fall Reduction in Chronic Obstructive Pulmonary Disease: Protocol for a Randomized Controlled Trial. *JMIR Res Protoc* 6, e228, doi:10.2196/resprot.8178 (2017).
- 249 Gonzalez Lindh, M. et al. Subjective swallowing symptoms and related risk factors in COPD. *ERJ Open Res* 5, doi:10.1183/23120541.00081-2019 (2019).
- 250 Good-Fratturelli, M. D., Curlee, R. F. & Holle, J. L. Prevalence and nature of dysphagia in VA patients with COPD referred for videofluoroscopic swallow examination. *J*

- Commun Disord 33, 93-110, doi:10.1016/s0021-9924(99)00026-x (2000).
- 251 Gross, R. D., Atwood, C. W., Jr., Ross, S. B., Olszewski, J. W. & Eichhorn, K. A. The coordination of breathing and swallowing in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 179, 559-565, doi:10.1164/rccm.200807-11390C (2009).
- 252 Loeb, M. B., Becker, M., Eady, A. & Walker-Dilks, C. Interventions to prevent aspiration pneumonia in older adults: a systematic review. *J Am Geriatr Soc* 51, 1018-1022, doi:10.1046/j.1365-2389.2003.51318.x (2003).
- 253 Ghannouchi, I., Speyer, R., Doma, K., Cordier, R. & Verin, E. Swallowing function and chronic respiratory diseases: Systematic review. *Respir Med* 117, 54-64, doi:10.1016/j.rmed.2016.05.024 (2016).
- 254 Zheng, Z. et al. Silent aspiration in patients with exacerbation of COPD. *Eur Respir J* 48, 570-573, doi:10.1183/13993003.00007-2016 (2016).
- 255 Regan, J., Lawson, S. & De Aguiar, V. The Eating Assessment Tool-10 Predicts Aspiration in Adults with Stable Chronic Obstructive Pulmonary Disease. *Dysphagia* 32, 714-720, doi:10.1007/s00455-017-9822-2 (2017).
- 256 Virvidaki, I. E., Nasios, G., Kosmidou, M., Giannopoulos, S. & Milionis, H. Swallowing and Aspiration Risk: A Critical Review of Non Instrumental Bedside Screening Tests. *J Clin Neurol* 14, 265-274, doi:10.3988/jcn.2018.14.3.265 (2018).
- 257 McKinstry, A., Tranter, M. & Sweeney, J. Outcomes of dysphagia intervention in a pulmonary rehabilitation program. *Dysphagia* 25, 104-111, doi:10.1007/s00455-009-9230-3 (2010).
- 258 Hussain, S. A. & Sullivan, R. Cancer control in Bangladesh. *Jpn J Clin Oncol* 43, 1159-1169, doi:10.1093/jjco/hyt140 (2013).
- 259 Li, Y. et al. Effect of emphysema on lung cancer risk in smokers: a computed tomography-based assessment. *Cancer Prev Res (Phila)* 4, 43-50, doi:10.1158/1940-6207.Ccrp-10-0151 (2011).
- 260 Dai, J., Yang, P., Cox, A. & Jiang, G. Lung cancer and chronic obstructive pulmonary disease: From a clinical perspective. *Oncotarget* 8, 18513-18524, doi:10.18632/oncotarget.14505 (2017).
- 261 Spyrtos, D., Papadaki, E., Lampaki, S. & Kontakiotis, T. Chronic obstructive pulmonary disease in patients with lung cancer: prevalence, impact and management challenges. *Lung Cancer (Auckl)* 8, 101-107, doi:10.2147/lctt.S117178 (2017).
- 262 Jankowich, M. D. & Rounds, S. I. S. Combined pulmonary fibrosis and emphysema syndrome: a review. *Chest* 141, 222-231, doi:10.1378/chest.11-1062 (2012).
- 263 Lin, H. & Jiang, S. Combined pulmonary fibrosis and emphysema (CPFE): an entity different from emphysema or pulmonary fibrosis alone. *J Thorac Dis* 7, 767-779, doi:10.3978/j.issn.2072-1439.2015.04.17 (2015).
- 264 Khan, R. et al. A New Variant of Combined Pulmonary Fibrosis and Emphysema From Second-Hand Smoke: A Case Report and Review of Literature. *J Clin Med Res* 7, 812-816, doi:10.14740/jocmr2277w (2015).
- 265 Papaioannou, A. I. et al. Combined pulmonary fibrosis and emphysema: The many aspects of a cohabitation contract. *Respir Med* 117, 14-26, doi:10.1016/j.rmed.2016.05.005 (2016).
- 266 Koo, H. J., Do, K.-H., Lee, J. B., Alblushi, S. & Lee, S. M. Lung cancer in combined pulmonary fibrosis and emphysema: a systematic review and meta-analysis. *PLoS one* 11, e0161437 (2016).
- 267 Crestani, B. et al. Long-term safety and tolerability of nintedanib in patients with idiopathic pulmonary fibrosis: results from the open-label extension study, INPULSIS-ON. *Lancet Respir Med* 7, 60-68, doi:10.1016/s2213-2600(18)30339-4 (2019).
- 268 Richeldi, L. et al. Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med* 370, 2071-2082, doi:10.1056/NEJMoa1402584 (2014).
- 269 Oltmanns, U. et al. Pirfenidone in idiopathic pulmonary fibrosis: real-life experience

- from a German tertiary referral center for interstitial lung diseases. *Respiration* 88, 199-207, doi:10.1159/000363064 (2014).
- 270 Serrano-Mollar, A. Cell therapy in idiopathic pulmonary fibrosis. *Medical sciences* 6, 64 (2018).
- 271 Rennard, S. I. & Daughton, D. M. Smoking cessation. *Clin Chest Med* 35, 165-176, doi:10.1016/j.ccm.2013.11.002 (2014).
- 272 Poole, P. J., Chacko, E., Wood-Baker, R. W. & Cates, C. J. Influenza vaccine for patients with chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*, Cd002733, doi:10.1002/14651858.CD002733.pub2 (2006).
- 273 Wongsurakiat, P. et al. Acute respiratory illness in patients with COPD and the effectiveness of influenza vaccination: a randomized controlled study. *Chest* 125, 2011-2020, doi:10.1378/chest.125.6.2011 (2004).
- 274 Nichol, K. L., Baken, L., Wuorenma, J. & Nelson, A. The health and economic benefits associated with pneumococcal vaccination of elderly persons with chronic lung disease. *Arch Intern Med* 159, 2437-2442, doi:10.1001/archinte.159.20.2437 (1999).
- 275 Nici, L., ZuWallack, R., Wouters, E. & Donner, C. (Eur Respiratory Soc, 2006).
- 276 Safka, K. A. & McIvor, R. A. Non-pharmacological management of chronic obstructive pulmonary disease. *Ulster Med J* 84, 13-21 (2015).
- 277 Spruit, M. A. et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med* 188, e13-64, doi:10.1164/rccm.201309-1634ST (2013).
- 278 Watz, H. et al. (Eur Respiratory Soc, 2014).
- 279 Hallal, P. C. et al. Global physical activity levels: surveillance progress, pitfalls, and prospects. *The lancet* 380, 247-257 (2012).
- 280 Organization, W. H. GPAQ global physical activity questionnaire. Accessed 2, 14 (2014).
- 281 Armstrong, T. & Bull, F. Development of the world health organization global physical activity questionnaire (GPAQ). *Journal of Public Health* 14, 66-70 (2006).
- 282 WHO. Physical Activity, <<https://www.who.int/news-room/fact-sheets/detail/physical-activity>> (2018).
- 283 Moniruzzaman, M. et al. Physical activity levels in Bangladeshi adults: results from STEPS survey 2010. *public health* 137, 131-138 (2016).
- 284 Jonkman, N. H. et al. Characteristics of effective self-management interventions in patients with COPD: individual patient data meta-analysis. *Eur Respir J* 48, 55-68, doi:10.1183/13993003.01860-2015 (2016).
- 285 NICE, G. U. T. U. Chronic obstructive pulmonary disease in over 16s: diagnosis and management. (2018).
- 286 Conti, G. et al. Noninvasive vs. conventional mechanical ventilation in patients with chronic obstructive pulmonary disease after failure of medical treatment in the ward: a randomized trial. *Intensive Care Med* 28, 1701-1707, doi:10.1007/s00134-002-1478-0 (2002).
- 287 Struik, F. M. et al. Nocturnal non-invasive ventilation in COPD patients with prolonged hypercapnia after ventilatory support for acute respiratory failure: a randomised, controlled, parallel-group study. *Thorax* 69, 826-834, doi:10.1136/thorax-jnl-2014-205126 (2014).
- 288 Marin, J. M., Soriano, J. B., Carrizo, S. J., Boldova, A. & Celli, B. R. Outcomes in patients with chronic obstructive pulmonary disease and obstructive sleep apnea: the overlap syndrome. *Am J Respir Crit Care Med* 182, 325-331, doi:10.1164/rccm.200912-1869OC (2010).
- 289 BTS. Clinical component for the home oxygen service in England and Wales, <[https://www.brit-thoracic.org.uk/document-library/clinical-information/oxygen/home-oxygen-guideline-\(adults\)/bts-home-oxygen-in-adults-clinical-component](https://www.brit-thoracic.org.uk/document-library/clinical-information/oxygen/home-oxygen-guideline-(adults)/bts-home-oxygen-in-adults-clinical-component)> (2006).

- 290 McNicholas, W. T., Verbraecken, J. & Marin, J. M. Sleep disorders in COPD: the forgotten dimension. *Eur Respir Rev* 22, 365-375, doi:10.1183/09059180.00003213 (2013).
- 291 Stoller, J. K., Panos, R. J., Krachman, S., Doherty, D. E. & Make, B. Oxygen therapy for patients with COPD: current evidence and the long-term oxygen treatment trial. *Chest* 138, 179-187, doi:10.1378/chest.09-2555 (2010).
- 292 Continuous or nocturnal oxygen therapy in hypoxemic chronic obstructive lung disease: a clinical trial. Nocturnal Oxygen Therapy Trial Group. *Ann Intern Med* 93, 391-398, doi:10.7326/0003-4819-93-3-391 (1980).
- 293 Ameer F, C. K., Usmani ZA, Smith BJ. Ambulatory oxygen for people with chronic obstructive pulmonary disease who are not hypoxaemic at rest (Review). *Cochrane Database of Systematic Reviews* (2014).
- 294 Roberts, C. M. Short burst oxygen therapy for relief of breathlessness in COPD. *Thorax* 59, 638-640, doi:10.1136/thx.2003.017301 (2004).
- 295 Quantrill, S. J. et al. Short burst oxygen therapy after activities of daily living in the home in chronic obstructive pulmonary disease. *Thorax* 62, 702-705, doi:10.1136/thx.2006.063636 (2007).
- 296 A family guide to ECMO, <<https://uihc.org/health-topics/family-guide-ecmo>> (2017).
- 297 Laros, C. D. et al. Bullectomy for giant bullae in emphysema. *J Thorac Cardiovasc Surg* 91, 63-70 (1986).
- 298 Murphy, P. B., Zoumot, Z. & Polkey, M. I. Noninvasive ventilation and lung volume reduction. *Clin Chest Med* 35, 251-269, doi:10.1016/j.ccm.2013.10.011 (2014).
- 299 Criner, G. J. et al. Effects of lung volume reduction surgery on gas exchange and breathing pattern during maximum exercise. *Chest* 135, 1268-1279, doi:10.1378/chest.08-1625 (2009).
- 300 Shah, P. L. et al. Endobronchial coils for the treatment of severe emphysema with hyperinflation (RESET): a randomised controlled trial. *Lancet Respir Med* 1, 233-240, doi:10.1016/s2213-2600(13)70047-x (2013).
- 301 Orens, J. B. et al. International guidelines for the selection of lung transplant candidates: 2006 update--a consensus report from the Pulmonary Scientific Council of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant* 25, 745-755, doi:10.1016/j.healun.2006.03.011 (2006).
- 302 Aziz, F., Penupolu, S., Xu, X. & He, J. Lung transplant in end-staged chronic obstructive pulmonary disease (COPD) patients: a concise review. *Journal of thoracic disease* 2, 111-116 (2010).
- 303 Spathis, A. & Booth, S. End of life care in chronic obstructive pulmonary disease: in search of a good death. *International journal of chronic obstructive pulmonary disease* 3, 11-29, doi:10.2147/copd.s698 (2008).
- 304 Gore, J. M., Brophy, C. J. & Greenstone, M. A. How well do we care for patients with end stage chronic obstructive pulmonary disease (COPD)? A comparison of palliative care and quality of life in COPD and lung cancer. *Thorax* 55, 1000-1006, doi:10.1136/thorax.55.12.1000 (2000).
- 305 Dean, M. M. End-of-life care for COPD patients. *Prim Care Respir J* 17, 46-50, doi:10.3132/pcrj.2008.00007 (2008).
- 306 Antus, B. Pharmacotherapy of Chronic Obstructive Pulmonary Disease: A Clinical Review. *ISRN Pulmonology* 2013, 582807, doi:10.1155/2013/582807 (2013).
- 307 Frith, P. A. et al. Efficacy and safety of the direct switch to indacaterol/glycopyrronium from salmeterol/fluticasone in non-frequently exacerbating COPD patients: The FLASH randomized controlled trial. *Respirology* 23, 1152-1159, doi:10.1111/resp.13374 (2018).
- 308 Lee, J. H., Kim, H. J. & Kim, Y. H. The Effectiveness of Anti-leukotriene Agents in

- Patients with COPD: A Systemic Review and Meta-analysis. *Lung* 193, 477-486, doi:10.1007/s00408-015-9743-5 (2015).
- 309 Rubinstein, I., Kumar, B. & Schriever, C. Long-term montelukast therapy in moderate to severe COPD--a preliminary observation. *Respir Med* 98, 134-138, doi:10.1016/j.rmed.2003.09.003 (2004).
- 310 Bateman, E. D. et al. Overall asthma control achieved with budesonide/formoterol maintenance and reliever therapy for patients on different treatment steps. *Respir Res* 12, 38, doi:10.1186/1465-9921-12-38 (2011).
- 311 Ishiura, Y. et al. Effect of triple therapy in patients with asthma-COPD overlap. *Int J Clin Pharmacol Ther* 57, 384-392, doi:10.5414/cp203382 (2019).
- 312 Bhattacharya, A. et al. Efficacy and safety of ipratropium bromide/salbutamol sulphate administered in a hydrofluoroalkane metered-dose inhaler for the treatment of COPD. *Int J Chron Obstruct Pulmon Dis* 11, 1469-1476, doi:10.2147/copd.S89923 (2016).
- 313 Cazzola, M., Calzetta, L., Rogliani, P., Page, C. & Matera, M. G. Impact of doxofylline in COPD: A pairwise meta-analysis. *Pulm Pharmacol Ther* 51, 1-9, doi:10.1016/j.pupt.2018.04.010 (2018).
- 314 Gaude, G. S. & Nadagouda, S. Nebulized corticosteroids in the management of acute exacerbation of COPD. *Lung India : official organ of Indian Chest Society* 27, 230-235, doi:10.4103/0970-2113.71957 (2010).
- 315 Ekström, T. & Tibblin, L. Influence of theophylline on gastro-oesophageal reflux and asthma. *Eur J Clin Pharmacol* 35, 353-356, doi:10.1007/bf00561363 (1988).
- 316 Dua, R. et al. Association of comorbid anxiety and depression with chronic obstructive pulmonary disease. *Lung India* 35, 31-36, doi:10.4103/lungindia.lungindia\_537\_16 (2018).
- 317 Kunik, M. E. et al. One session cognitive behavioural therapy for elderly patients with chronic obstructive pulmonary disease. *Psychol Med* 31, 717-723, doi:10.1017/s0033291701003890 (2001).
- 318 Giardino, N. D. et al. Anxiety is associated with diminished exercise performance and quality of life in severe emphysema: a cross-sectional study. *Respir Res* 11, 29, doi:10.1186/1465-9921-11-29 (2010).
- 319 Smoller, J. W., Pollack, M. H., Otto, M. W., Rosenbaum, J. F. & Kradin, R. L. Panic anxiety, dyspnea, and respiratory disease. Theoretical and clinical considerations. *Am J Respir Crit Care Med* 154, 6-17, doi:10.1164/ajrccm.154.1.8680700 (1996).
- 320 Cafarella, P. A., Effing, T. W., Usmani, Z. A. & Frith, P. A. Treatments for anxiety and depression in patients with chronic obstructive pulmonary disease: a literature review. *Respirology* 17, 627-638, doi:10.1111/j.1440-1843.2012.02148.x (2012).
- 321 Vozoris, N. T. et al. Benzodiazepine drug use and adverse respiratory outcomes among older adults with COPD. *Eur Respir J* 44, 332-340, doi:10.1183/09031936.00008014 (2014).
- 322 Vozoris, N. T. et al. Benzodiazepine use among older adults with chronic obstructive pulmonary disease: a population-based cohort study. *Drugs Aging* 30, 183-192, doi:10.1007/s40266-013-0056-1 (2013).
- 323 Zimlich, R. Benzodiazepine use in older COPD patients may not be ideal. *Medical Economics* 15 (2015).
- 324 Battaglia, S., Bezzi, M. & Sferrazza Papa, G. F. Are benzodiazepines and opioids really safe in patients with severe COPD? *Minerva Med* 105, 1-7 (2014).
- 325 Hunter, M. H. & King, D. E. COPD: management of acute exacerbations and chronic stable disease. *Am Fam Physician* 64, 603-612 (2001).
- 326 Aloush, V., Navon-Venezia, S., Seigman-Igra, Y., Cabili, S. & Carmeli, Y. Multidrug-resistant *Pseudomonas aeruginosa*: risk factors and clinical impact. *Antimicrob Agents Chemother* 50, 43-48, doi:10.1128/aac.50.1.43-48.2006 (2006).



- 327 Montero, M. et al. Risk factors for multidrug-resistant *Pseudomonas aeruginosa* acquisition. Impact of antibiotic use in a double case-control study. *Eur J Clin Microbiol Infect Dis* 29, 335-339, doi:10.1007/s10096-009-0850-1 (2010).
- 328 Macfarlane, J. T., Colville, A., Guion, A., Macfarlane, R. M. & Rose, D. H. Prospective study of aetiology and outcome of adult lower-respiratory-tract infections in the community. *Lancet* 341, 511-514, doi:10.1016/0140-6736(93)90275-I (1993).
- 329 Patel, I. S. et al. Relationship between bacterial colonisation and the frequency, character, and severity of COPD exacerbations. *Thorax* 57, 759-764, doi:10.1136/thorax.57.9.759 (2002).
- 330 Murphy, T. F., Sethi, S., Klingman, K. L., Brueggemann, A. B. & Doern, G. V. Simultaneous respiratory tract colonization by multiple strains of nontypeable haemophilus influenzae in chronic obstructive pulmonary disease: implications for antibiotic therapy. *J Infect Dis* 180, 404-409, doi:10.1086/314870 (1999).
- 331 Mogulkoc, N. et al. Acute purulent exacerbation of chronic obstructive pulmonary disease and Chlamydia pneumoniae infection. *Am J Respir Crit Care Med* 160, 349-353, doi:10.1164/ajrccm.160.1.9809041 (1999).
- 332 Soler, N. et al. Bronchial microbial patterns in severe exacerbations of chronic obstructive pulmonary disease (COPD) requiring mechanical ventilation. *Am J Respir Crit Care Med* 157, 1498-1505, doi:10.1164/ajrccm.157.5.9711044 (1998).
- 333 Montero, M., Domínguez, M., Orozco-Levi, M., Salvadó, M. & Knobel, H. Mortality of COPD patients infected with multi-resistant *Pseudomonas aeruginosa*: a case and control study. *Infection* 37, 16-19, doi:10.1007/s15010-008-8125-9 (2009).
- 334 Butler, C. C. et al. C-Reactive Protein Testing to Guide Antibiotic Prescribing for COPD Exacerbations. *N Engl J Med* 381, 111-120, doi:10.1056/NEJMoa1803185 (2019).
- 335 Food, U. & Administration, D. Guidance for industry and FDA staff: Review criteria for assessment of C-reactive protein (CRP), high sensitivity C-reactive protein (hsCRP), and cardiac C-reactive protein (cCRP) assays. Silver Spring, MD: US Center for Devices and Radiological Health (2005).
- 336 Gilbert, D. N. Procalcitonin as a biomarker in respiratory tract infection. *Clin Infect Dis* 52 Suppl 4, S346-350, doi:10.1093/cid/cir050 (2011).
- 337 Schuetz, P., Albrich, W. & Mueller, B. Procalcitonin for diagnosis of infection and guide to antibiotic decisions: past, present and future. *BMC Med* 9, 107, doi:10.1186/1741-7015-9-107 (2011).
- 338 Daniels, J. M. et al. Antibiotics in addition to systemic corticosteroids for acute exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 181, 150-157, doi:10.1164/rccm.200906-0837OC (2010).
- 339 Rello, J. Importance of appropriate initial antibiotic therapy and de-escalation in the treatment of nosocomial pneumonia. *European Respiratory Review* 16, 33-39 (2007).
- 340 Micek, S. T., Kollef, K. E., Reichley, R. M., Roubinian, N. & Kollef, M. H. Health care-associated pneumonia and community-acquired pneumonia: a single-center experience. *Antimicrob Agents Chemother* 51, 3568-3573, doi:10.1128/aac.00851-07 (2007).
- 341 Cates, C. Spacers and nebulisers for the delivery of beta-agonists in non-life-threatening acute asthma. *Respir Med* 97, 762-769, doi:10.1016/s0954-6111(03)00021-0 (2003).
- 342 Kopsaftis, Z. A. et al. Short-acting bronchodilators for the management of acute exacerbations of chronic obstructive pulmonary disease in the hospital setting: systematic review. *Syst Rev* 7, 213, doi:10.1186/s13643-018-0860-0 (2018).
- 343 Adkison, J. D. & Konzem, S. L. Management of acute exacerbations of chronic obstructive pulmonary disease. *Pharmacotherapy* 21, 929-939, doi:10.1592/phco.21.11.929.34523 (2001).
- 344 Walters, J. A., Tan, D. J., White, C. J. & Wood-Baker, R. Different durations of cor-

- ticosteroid therapy for exacerbations of chronic obstructive pulmonary disease. Cochrane Database Syst Rev, Cd006897, doi:10.1002/14651858.CD006897.pub3 (2014).
- 345 Gupta, H. et al. Impact of COPD on postoperative outcomes: results from a national  
database. *Chest* 143, 1599-1606, doi:10.1378/chest.12-1499 (2013).
- 346 Fields, A. C. & Divino, C. M. Surgical outcomes in patients with chronic obstructive  
pulmonary disease undergoing abdominal operations: An analysis of 331,425 pa-  
347 tients. *Surgery* 159, 1210-1216, doi:10.1016/j.surg.2015.11.007 (2016).
- Cukic, V. Preoperative prediction of lung function in pneumonectomy by spirome-  
348 try and lung perfusion scintigraphy. *Acta Inform Med* 20, 221-225, doi:10.5455/  
aim.2012.20.221-225 (2012).
- Oswald, N. K. et al. Predicting Postoperative Lung Function Following Lung Can-  
349 cer Resection: A Systematic Review and Meta-analysis. *EClinicalMedicine* 15, 7-13,  
doi:10.1016/j.eclinm.2019.08.015 (2019).
- Balkissoon, R. Stem Cell Therapy for COPD: Where are we? *Chronic Obstr Pulm Dis* 5,  
350 148-153, doi:10.15326/jcopdf.5.2.2018.0140 (2018).
- Han, C. et al. Exosomes and Their Therapeutic Potentials of Stem Cells. *Stem Cells Int*  
351 2016, 7653489-7653489, doi:10.1155/2016/7653489 (2016).
- Ratajczak, M. Z., Jadczyk, T., Pędziwiatr, D. & Wojakowski, W. New advances in stem  
cell research: practical implications for regenerative medicine. *Pol Arch Med Wewn*  
352 124, 417-426, doi:10.20452/pamw.2355 (2014).
- O'Farrell, H. E. & Yang, I. A. Extracellular vesicles in chronic obstructive pulmonary  
disease (COPD). *J Thorac Dis* 11, S2141-s2154, doi:10.21037/jtd.2019.10.16  
(2019).

# BANGLADESH GUIDELINES FOR CHRONIC OBSTRUCTIVE PULMONARY DISEASE

## 2020

Endorsed By:

- Bangladesh Primary Care Respiratory Society



Designed and Developed by:  
Ingenious Health Care Limited  
41 Ring Road (2nd Floor), Mohammadpur, Dhaka-1207



*Printing Courtesy by:*

